

A quantitative study of modern pollen-vegetation relationships in southern Brazil's Araucaria forest

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Cardenas, M., Wilson, O., Schorn, L., Mayle, F. and Iriarte, J. (2019) A quantitative study of modern pollen-vegetation relationships in southern Brazil's Araucaria forest. *Review of Palaeobotany and Palynology*, 265. pp. 27-40. ISSN 0034-6667 doi: <https://doi.org/10.1016/j.revpalbo.2019.03.003> Available at <https://centaur.reading.ac.uk/83038/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.revpalbo.2019.03.003>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 A quantitative study of modern pollen-
2 vegetation relationships in southern Brazil's
3 *Araucaria* forest.

4 **Macarena L. Cárdenas^{a,b}, Oliver J. Wilson^{a*}, Lauri A. Schorn^c, Francis E. Mayle^a, José Iriarte^d**

5 ^a School of Archaeology, Geography and Environmental Science (SAGES), University of Reading,
6 Whiteknights, P.O. Box 227, Reading RG6 6DW, Berkshire, United Kingdom

7 ^b Present address: Earthwatch Institute, Mayfield house, 256 Banbury Road, Oxford, OX2 7DE,
8 United Kingdom

9 ^c Departamento e Engenharia Florestal, Universidade Regional de Blumenau, Brazil, 47-3221-6043,
10 47-99973-4820

11 ^d Department of Archaeology, University of Exeter, North Park Rd, Exeter, EX4 4QE, United Kingdom

12 * Corresponding author: o.j.wilson@reading.ac.uk

13 **Abstract**

14 Southern Brazil's highland *Araucaria* forest is ancient, diverse and unique, but its future is under
15 significant threat from 20th Century habitat loss and 21st Century climate change. Palaeoecological
16 studies have revealed that it expanded rapidly over highland grasslands around 1,000 years ago,
17 but whether this expansion was caused by human land use or climate change has been a topic of
18 considerable debate. Discriminating between these potential drivers has so far not been possible

19 with fossil pollen, however, as the palynological representation of floristic and structural
20 differences in *Araucaria* forest remains poorly understood. Here, we address this shortcoming using
21 modern pollen rain from moss polsters and vegetation surveys in forest areas with minimal current
22 human disturbance. We show that forest plots with evident structural differences lack consistent
23 differences in their floristic composition and cannot be reliably distinguished by their pollen
24 spectra. We quantify pollen-vegetation relationships for 27 key tree genera of *Araucaria* forest,
25 showing that, despite significant intra-taxon variability, 22 of these are under-represented or
26 absent in the pollen record. These palynologically under-represented and silent taxa include many
27 of the forest's most ecologically important tree species, with only *Araucaria*, *Lamanonia*,
28 *Podocarpus*, *Myrsine* and *Clethra* being more abundant in the pollen rain than vegetation. Our
29 results suggest that subtle structural changes in *Araucaria* forest, as well as moderate to significant
30 floristic changes, may not be clearly distinguished in fossil pollen records – an important limitation
31 when attempting to identify past human and climatic impacts on *Araucaria* forest via pollen
32 analysis.

33 Key words: *Araucaria* forest, pollen, moss polsters, modern analogues, human impact

34 1. Introduction

35 1.1. Rationale

36 Disentangling the past effects of humans and climate on ecosystems requires a rigorous
37 understanding of the vegetation changes that occurred, and how these are reflected in
38 palaeoecological proxies. This is an important undertaking in Brazil's *Araucaria* forest (also known
39 as Mixed Ombrophilous Forest), an ancient, highly diverse and threatened ecosystem in the
40 country's southern highlands. Studies have charted changes in the distribution of *Araucaria* forest
41 in relation to climatic variations since the late Pleistocene (Behling et al., 2004; Ledru et al., 2009),

42 many finding rapid increases in its pollen abundance about 1,000 years ago which, in some sites,
43 reached levels unprecedented for tens of thousands of years (Behling et al., 2004; Gu et al., 2017).
44 This marked expansion ca. 1000 years ago was much more drastic than the steady expansion of
45 *Araucaria* forest over the previous few millennia, and has been attributed by some authors to
46 ancient human land use, rather than climate change, due to the coincident expansion of the local
47 southern proto-Jê culture (Bitencourt and Krauspenhar, 2006; dos Reis et al., 2014; Lauterjung et
48 al., 2018; Noelli, 2000; Robinson et al., 2018), whose economy, spiritual beliefs and living space
49 were closely linked to the forest (De Souza et al., 2016a; Iriarte et al., 2016; Iriarte and Behling,
50 2007).

51 Understanding the extent to which *Araucaria* forest's late-Holocene expansion can be attributed to
52 humans versus climate change will improve our understanding of the ecosystem's responses to
53 prevailing climatic conditions – a particularly important endeavour in the face of continuing
54 anthropogenic climate change. With regional temperatures forecast to become warmer, and
55 precipitation more variable, as this century progresses (Chou et al., 2014), the cold- and humidity-
56 dependent *Araucaria* forest is likely to contract (Bergamin, 2017; Wrege et al., 2009).

57 Understanding how this iconic forest responded to past variations in climate – and how humans
58 may have altered this response – may help conservationists anticipate and ameliorate the effects of
59 modern climatic changes upon its distribution. This is especially important as *Araucaria* forest cover
60 was much reduced by the 20th Century logging boom in the states of Paraná, Santa Catarina and
61 Rio Grande do Sul (Carvalho and Nodari, 2010), with remnants now covering just 12.6% of the
62 biogeographic region it previously dominated (Ribeiro et al., 2009).

63 The main proxy used for reconstructing past dynamics of vegetation on the southern Brazilian
64 highlands is fossil pollen from lake/bog sediments, the informed and accurate interpretation of

65 which requires appropriate characterisation and quantification of modern pollen-vegetation
66 relationships. This is currently poor for Brazil's *Araucaria* forest. Existing pollen-vegetation studies
67 (Behling et al., 2001; Garcia et al., 2004; Jeske-Pieruschka et al., 2010) are purely qualitative,
68 identifying *Araucaria* forest's key indicator taxa but giving little insight into the degree to which the
69 floristic composition or structure of these forests can be resolved palynologically. In this study, we
70 address this shortcoming, presenting the first quantitative examination of the modern pollen-
71 vegetation relationship of *Araucaria* forest and its constituent taxa.

72 1.2. Background

73 *Araucaria* forest is found at the southern and south-eastern reaches of the Atlantic Forest biome, a
74 global biodiversity hotspot (Colombo and Joly, 2010; Myers et al., 2000). It occupies the biome's
75 high-altitude and low-temperature extreme niches (Neves et al., 2017; Oliveira-Filho et al., 2014),
76 mostly occurring above 500m altitude (Hueck, 1953; Lacerda, 2016) in areas with high annual rainfall
77 and climatic conditions classified as Cfb – humid subtropical oceanic climate, with temperate
78 summers and no dry season (Alvares et al., 2013) – under the Köppen system (Higuchi et al., 2012;
79 Hueck, 1953). At more southerly latitudes in Brazil, some apparently natural stands of *Araucaria*
80 forest can also be found at elevations below 500m (Behling et al., 2016).

81 *Araucaria* forest is home to over 1,500 species of plants, 6% of which are endemic to the Atlantic
82 Forest biome (Neves et al., 2017), and is characterised by the dominance of the conifer *Araucaria*
83 *angustifolia* in its canopy (Duarte et al., 2014; Leite and Klein, 1990; Meyer et al., 2013). This
84 species has the third highest EDGE score (a metric which combines evolutionary distinctiveness and
85 global endangerment) of all the world's gymnosperms (Forest et al., 2018), imperilled by logging
86 and habitat loss which started in the late 19th Century and reduced the forests' area by 97% within
87 three generations (Carvalho and Nodari, 2010; Thomas, 2013). Other important arboreal species

88 include *Dicksonia sellowiana*, *Podocarpus sellowiana*, *Matayba elaeagnoides*, *Lithraea brasiliensis*,
89 *Clethra scabra*, *Ocotea porosa*, and *Prunus myrtifolia* (Duarte et al., 2014; Meyer et al., 2013).

90 *Araucaria* forest is not floristically homogenous, however. Early studies differentiated eight types of
91 *Araucaria* forest (Klein, 1978), and the ecosystem's mixture of tropical and temperate elements
92 varies with longitude, altitude, temperature, river basin, and distance to other forest types (Duarte
93 et al., 2014; Gonçalves and Souza, 2014; Higuchi et al., 2012; Klein, 1975; Oliveira-Filho et al., 2014;
94 Sevegnani et al., 2016). At higher altitudes, under drier conditions and in areas of high fire
95 frequencies and shallower soils, *Araucaria* forest forms mosaics with campos grasslands (Hueck,
96 1966; Müller et al., 2012; Oliveira-Filho et al., 2014), with other ecotones occurring with seasonally
97 deciduous and dense ombrophilous forests at its western and eastern extents, respectively
98 (Gonçalves and Souza, 2014; Oliveira-Filho et al., 2014).

99 Fossil pollen records from the eastern edge of southern Brazil's highland plateau show that the area
100 was dominated by campos at the Last Glacial Maximum, with *Araucaria* forest likely confined to
101 river valleys (Behling, 2002; Behling et al., 2004; Leonhardt and Lorscheitter, 2010). These forests
102 subsequently expanded slightly 4,000-3,000 years ago, and then rapidly 1,500-900 years ago
103 (Behling, 1997a, 1995; Behling et al., 2004; Iriarte and Behling, 2007; Jeske-Pieruschka and Behling,
104 2012; Leonhardt and Lorscheitter, 2010; Silva and Anand, 2011), in some places replacing
105 grasslands within a century (Behling et al., 2004; Iriarte and Behling, 2007).

106 These changes have typically been attributed to climate change (Rodrigues et al., 2016a). However,
107 recent speleothem data (Bernal et al., 2016) show that, although the initial *Araucaria* forest
108 expansion at 4,000 yr BP does correlate with an increase in rainfall, no subsequent climate change
109 accompanies the later, more significant forest expansion at 1,500-900 yr BP (Robinson et al., 2018).
110 This has led several authors to invoke humans as the most likely cause for this most recent, rapid

111 expansion of *Araucaria* forest (Bitencourt and Krauspenhar, 2006; dos Reis et al., 2014; Iriarte and
112 Behling, 2007; Noelli, 2000), bolstered by evidence that, under natural conditions, expansion of
113 forest patches into campos grassland can be extremely slow (less than 100m in 4,000 years (Silva
114 and Anand, 2011)).

115 Before European arrival, the indigenous people of the highlands, the southern Jê, had lifestyles that
116 were at least semi-sedentary and practised a mixed economy that combined the cultivation of
117 manioc (*Manihot esculenta*), maize (*Zea mays*), squash (*Cucurbita* sp.) and beans (*Phaseolus* sp.)
118 with hunting, fishing, and gathering forest and riverine resources (Corteletti et al., 2015; De Souza
119 et al., 2016b; Henry, 1964; Métraux, 1946; Noelli and De Souza, 2017). *Araucaria angustifolia* seeds,
120 known as pinhão, have long been a critical food source for the southern Jê (Corteletti et al., 2015;
121 Heineberg, 2014; Henry, 1964; Loponte et al., 2016; Métraux, 1946; Urban, 1985), with one group
122 traditionally defining a year as “one period with no pine nuts” (Henry, 1964, p. 68). Ethnographic
123 studies of extant southern Jê groups have also highlighted the importance of *Araucaria* forest – and
124 *A. angustifolia* in particular – as a source of spiritual power and cultural identity (Fernandes and
125 Piovezana, 2015; Haverroth, 1997; Hoffmann, 2011; Robinson et al., 2017; Silva, 2002).

126 Archaeological records show that the southern Jê flourished around 1,000 years ago (Iriarte et al.,
127 2016, 2013; Noelli and De Souza, 2017; Robinson et al., 2018), as changes in domestic architecture
128 appeared (De Souza et al., 2016b), combining with the arrival of funerary mound-and-enclosure
129 complexes (De Souza et al., 2016a; Iriarte et al., 2016) to form highly structured ‘sacred landscapes’
130 (Iriarte et al., 2013) in the highlands. The temporal overlap of these changes with the most recent
131 *Araucaria* forest expansion suggests that more available forest resources likely helped underpin the
132 cultural changes (Iriarte and Behling, 2007), and there is some evidence that the southern Jê may in

turn have helped expand the forest (dos Reis et al., 2014; Lauterjung et al., 2018; Robinson et al., 2018), though this has not yet been tested with fossil pollen data.

1.3. Aims

Quantitative studies of the relationships between present-day *Araucaria* forest and its constituent pollen assemblages are needed to improve the level of ecological detail that can be extracted from fossil pollen records, and thereby obtain a firmer basis for unravelling the relative roles of climate change versus human land use in driving this threatened forest's long-term dynamics over the past several millennia. We therefore conducted a study to quantitatively define the vegetation-pollen relationships of *Araucaria* forest taxa. Specifically, we analysed floristic inventories from vegetation plots, and collected their constituent pollen assemblages from moss polsters, to determine: 1. whether structurally different *Araucaria* forest patches can be differentiated by their pollen spectra, and 2. how the constituent species of these patches are represented in the modern pollen rain.

2. Methods

2.1. Study site

[Insert Figure 1 here]

Figure 1: Map of forest plot locations; numbers refer to plot codes (see Table 1 below). Plot colours: Cyan: riverine; Olive: slope; Black: rock outcrop; Purple: open understorey; Orange: closed understorey; Red: disturbed.

The Reserva Particular do Patrimônio Natural (RPPN) Emilio Einsfeld Filho (27°58'0.80"S, 50°49'20.03"W, 650-990 m a.s.l.), managed by the Florestal Gateados company, is located in the municipalities of Campo Belo do Sul and Capão Alto, southern Santa Catarina State, Brazil (Figure

155 1). RPPN Emilio Einsfeld Filho is located on the plateau of the Serra Catarinense, a gently undulating
156 landscape. The reserve covers 6,329 hectares of native *Araucaria* forest patches within a wider
157 matrix of campos grassland, with the largest density of forest along rivers, valleys and slopes.
158 Forested areas have been protected from timber extraction since 1989 and cattle grazing since
159 1993 (Zeller, 2010).

160 The climate of the area corresponds to Cfb in the Köppen classification: humid subtropical with
161 temperate summers and no dry season, but with occurrence of severe frosts (Alvares et al., 2013;
162 Formento et al., 2004; Zeller, 2010). Between 2005 and 2009 the reserve's average annual
163 temperature was 16°C, with hottest months averaging 31°C, coldest months averaging 6.5°C and a
164 lowest recorded temperature of -12°C; in this period annual precipitation varied between 1300 and
165 2400 mm, with an average of 129 rainy days a year (Zeller, 2010).

166 2.2 Field sampling

167 2.2.1 Vegetation survey

168 The native *Araucaria* forest at RPPN Emilio Einsfeld Filho has been subject to ecological studies for
169 the last 20 years, with permanent plots of 10 x 50 m installed to perform floristic inventories and
170 study forest dynamics under distinct geographic conditions. Within each plot all trees with d.b.h.
171 (diameter at breast height) ≥ 10 cm have been recorded (Formento et al., 2004; Maçaneiro et al.,
172 2018). Percentage abundances were calculated based upon all recorded stems surveyed within
173 each plot.

174 The sixteen plots used in this study are located within the oldest and least disturbed areas of forest
175 and incorporate the widest available range of vegetation form and physical setting (Table 1). Data
176 on canopy openness, slope, animal disturbance, geographical feature associations and topographic
177 position were collected during botanical surveys in 2015, and the plots classified into six structural

178 categories according to their dominant feature:

- 179
- slope: forest patches in areas with > 25% inclination
- 180
- rock outcrop: forest patches where rock outcrops cover > 20% of plot surface
- 181
- riverine: forest patch bounded by, or entirely within 50 m of, a water course (river or creek)
- 182
- open understorey: forest with closed canopy and scarce understorey
- 183
- closed understorey: forest with closed canopy and dense understorey
- 184
- naturally disturbed: forest patches with evidence of ground disturbance by wild boars

Plot ID	Forest Plot category	Coordinates	Altitude (a.s.l.)
114	Slope	28° 2'34.70"S, 50°45'21.40"W	780 m
118	Slope	28° 3'28.80"S, 50°45'45.80"W	924 m
117	Rock outcrop	28° 3'0.10"S, 50°45'16.50"W	856 m
110	Riverine	28° 1'17.30"S, 50°46'55.50"W	721 m
136	Riverine	28° 1'55.50"S, 50°45'59.00"W	740 m
151	Riverine	28° 3'22.50"S,	874 m

		50°45'29.60"W	
119	Open understorey	28° 3'16.80"S, 50°45'51.70"W	929 m
122	Open understorey	28° 2'59.00"S, 50°46'28.10"W	921 m
123	Open understorey	28° 3'4.10"S, 50°46'11.60"W	935 m
142	Open understorey	28° 3'42.60"S, 50°46'26.60"W	922 m
144	Open understorey	28° 3'24.30"S, 50°46'21.50"W	953 m
156	Disturbed	28° 4'29.00"S, 50°46'22.10"W	850 m
157	Disturbed	28° 4'11.80"S, 50°46'20.90"W	851 m
159	Disturbed	28° 4'21.60"S, 50°45'53.60"W	868 m
141	Closed understorey	28° 3'40.70"S, 50°46'14.90"W	931 m
149	Closed understorey	28° 3'53.50"S, 50°45'39.20"W	891 m

185 Table 1: The studied forest plots, their structural category, and the location and altitude of each
186 transect's starting point.

187 2.2.2 Modern pollen rain

188 Using moss polsters to study modern pollen rain has been widely used and recognized as a reliable
189 technique (Atanassova, 2007; Caseldine, 1989; Hicks, 1977; López-Sáez et al., 2010; Pardoe, 2006;
190 Pardoe et al., 2010; Tonkov et al., 2001). Moss polsters can collect several years' pollen (Pardoe et
191 al., 2010), occur naturally within the forest on different substrates and at varied heights (allowing
192 for effective capture of pollen rain) and do not need to be installed and managed before analysis
193 (Hicks, 1985). Moss polsters were gathered every 10 m along the 50 m length of each forest
194 inventory plot surveyed. At each collection point, polsters were obtained from ground level
195 (including rocks), chest height (1.20 m) and overhead (1.80 m). The samples for each point of
196 collection were mixed to form one homogeneous sample for each plot (following Pardoe et al.,
197 2010), placed in a labelled, sealed plastic bag, and stored at 4°C to preserve the material.

198 2.3. Laboratory methods

199 Subsamples of 1 cm³ were extracted from each homogenised sample for pollen processing. Each
200 subsample was centrifuged, washed with distilled water, centrifuged again, then treated with 10%
201 NaOH and put in a hot-water bath to deflocculate the organics. Afterwards, acetolysis was carried
202 out in the samples to dissolve cellulose material (Faegri and Iversen, 1989).

203 Samples were mounted on slides in silicone oil and counted at 400× and 1000× magnification using
204 a Leica DME binocular microscope. Samples were counted to a minimum of 300 terrestrial pollen
205 grains. Aquatic taxa and spores were also counted as is standard practice, but were excluded from
206 the total terrestrial pollen (TTP) sum; non-native species (e.g. pine from recent plantations
207 surrounding the reserve) were also excluded from the TTP sum. For pollen determination, the

reference collection of the Tropical Palaeoecology Research Group of the University of Reading was used, along with several pollen atlases – Colinvaux et al. (1999), Hooghiemstra (1984), Roubik and Moreno (1991), Behling (1993) – and the online Neotropical Pollen Key (Bush and Weng, 2007). Diagrams of vegetation and pollen abundance were made using C2 v.1.7.7 (Juggins, 2016).

2.4. Data analysis

2.4.1 Vegetation data processing

Analysis of each plot's floristic inventory data used percentage abundance (relative to the total number of individual trees counted) and relative coverage value (RCV). RCV (expressed as a percentage) indicates the importance of a species within a forest plot, taking into account its individuals' density and dominance.

$$RCVi = \frac{(RDoi + RDeni)}{\sum(RDo + RDen)} * 100$$

RCVi: Relative Coverage Value for species *i* (%)

RDoi: Relative dominance for species *i* (basal area (m²) per hectare)

RDeni: Relative density for species *i* (number of individuals per hectare)

RDo: Relative dominance for all species (m² / ha)

RDen: Relative density for all species (n / ha)

2.4.2 Statistical analysis of vegetation and pollen data

To investigate the extent to which structural differences between plots are reflected in their vegetation composition and pollen rain, we conducted cluster and Detrended Correspondence Analysis (DCA) multivariate analyses using PAST (Hammer et al., 2001) v.3.19 (2018). Raw floristic inventory and pollen data were normalised using natural logarithm before running the analyses. Only taxa with abundance ≥ 2% in two or more plots were considered for cluster and multivariate

230 analysis.

231 Cluster analysis was run with hierarchical clustering using the Unweighted Pair Group Method
232 (UPGM) algorithm, which iteratively joins clusters based on the average distance between all
233 members of the groups (Kent and Coker, 1992; Legendre et al., 2012). The measure of similarity or
234 distance measurements between samples (forest plots) was calculated using Bray-Curtis index of
235 dissimilarity (Kent and Coker, 1992). The DCA uses the same algorithm as Decorana (Hill and Gauch,
236 1980), with modifications according to Oksanen and Minchin (1997). The DCA method was
237 preferred as it allows the identification of the ecological variances amongst the forest plots (Correa-
238 Metrio et al., 2014).

239 To enable direct comparisons between taxa in the vegetation and pollen rain, which were mostly
240 identified at different taxonomic levels, the vegetation-pollen analysis was performed at genus
241 level, with congeneric species combined for the analysis (Burn et al., 2010).

242 To quantify the vegetation-pollen relationships of taxa in this study area, we calculated mean p/v
243 (pollen/vegetation) values (Burn et al., 2010; Gosling et al., 2009; Montade et al., 2016), which are
244 analogous to R values (Davis, 1963). We calculate the p/v value for taxon i as:

245
$$p/v_i = \frac{\text{Pollen abundance (\%) averaged over all plots}_i}{\text{Vegetation abundance (\%) averaged over all plots}_i}$$

246 Although more complex measures of pollen production exist (e.g. Parsons and Prentice, 1981;
247 Sugita, 2007a, 2007b), p/v values provide a straightforward and intuitive index of the relationship
248 between vegetation and pollen abundance and continue to be foundational for vegetation
249 reconstruction (Mrotzek et al., 2017).

250 To complement this metric and show the variability in pollen-vegetation relationships between
251 plots, we also calculated a pollen-vegetation abundance differential for taxon i :

$$Abundance\ differential_i = Pollen\ abundance\ (\%)_i - Vegetation\ abundance\ (\%)_i$$

Plots where a taxon was absent from both the pollen rain and vegetation survey were not included in calculations of mean differentials. p/v values and abundance differentials could not be calculated for herbaceous taxa as these were not included in the vegetation survey. Data were plotted using R v.3.4.2 (R Core Team, 2017) and PAST (Hammer et al., 2001) v.3.19 (2018).

3. Results

3.1. Vegetation

Across the 16 forest plots, 54 tree species with d.b.h. ≥ 10 cm were recorded, of which 33 had relative abundances of 2% or higher in at least two plots; the relative (percentage) abundance of these species within each plot is shown in Figure 2. The most consistently present species were *Lithraea brasiliensis* (15 plots), *Cinnamodendron dinisii* (13 plots), *Myrsine coriacea*, *Ocotea pulchella* and *Araucaria angustifolia* (12 plots), undifferentiated *Eugenia* sp. (11 plots), and *Calypttranthes* cf. *concinna*, *Ilex theezans*, *Pera glabrata*, *Podocarpus lambertii* and *Zanthoxylum kleinii* (10 plots).

The taxa with the highest average abundance in the vegetation were *L. brasiliensis* (9.1%), *Cinnamodendron dinisii* (7.7%), *A. angustifolia* (7.3%), undifferentiated *Eugenia* sp. (6.7%), *O. pulchella* (5.7%), and *Calypttranthes* cf. *concinna* (5.0%). Of the common taxa, *L. brasiliensis* had the highest maximum abundance (34.2%), followed by *Eugenia* sp. (26.4%), *A. angustifolia* (26.1%), *E. uniflora* (25.7%) and *Cinnamodendron dinisii* (24.1%). Two less common species with high maximum abundances were *Matayba elaeagnoides* (found in three plots, max. 35.6%) and *Nectandra megapotamica* (three plots, max. 28.6%).

Generally, tree species that were more abundant also had higher relative coverage values (RCVs;

Figure 3): the taxa with the highest RCVs averaged across all plots were *L. brasiliensis* (10.3%), *A. angustifolia* (10.0%), *O. pulchella* (7.8%), *C. dinisii* (6.1%), *Styrax leprosus* (5.3%) and undifferentiated *Eugenia* sp. (5.2%). Some species were found in few plots, but had high RCVs where they were present: the less frequent species with the highest RCVs averaged across the plots in which they were found were *M. elaeagnoides* (15.6%, four plots), *Luehea divaricata* (9.2%, three plots), *N. megapotamica* (9.0%, three plots), *Vernonanthura discolor* (7.4%, five plots), *Sebastiania commersoniana* (7.0%, six plots), *Prunus myrtifolia* (5.7%, six plots), *Piptocarpha angustifolia* (5.2%, two plots), *Myrceugenia* sp. (5.1%, five plots) and *Myrsine umbellata* (5.0%, two plots).

3.2. Pollen

170 pollen and spore types were found in the 16 forest plots. Of these, 118 could be identified to family level, 95 of which could be refined to genus, one to species (*Ilex theezans*). We also assign species names to *Araucaria angustifolia* and *Podocarpus lambertii* because no other species of these genera occur in our study area. The plots' pollen spectra are summarised in Figure 4, where abundant and important taxa are shown.

Eight pollen types were found in all 16 plots: *A. angustifolia*, Asteraceae (others total), *Eugenia* type, 'cf *Lithraea* a (Anacardiaceae)', *Myrsine*, Poaceae, *P. lambertii*, and Pteridophyta. Additionally, Araceae pollen was found in 15 plots, and *Lithraea*, *Vernonanthura*-type and *Calyptanthus*-type in 14; *Sebastiania* and undifferentiated Myrtaceae pollen was found in 12 plots, and *Clethra*, *I. theezans* and *Pteris* each in 10 plots. Additionally, *Alchornea* pollen and *Cyathea*-type spores were found at very low levels (average abundance 0.27% and 0.2% respectively) in 11 plots, with *Senecio*-type (average 0.35%) and *Mimosa scabrella*-type (average 0.25%) present in 10 plots. *Myrsine* pollen was the most abundant across all plots (averaging 19.5%), followed by *A. angustifolia* (16.6%), Pteridophyta (13.2%), *Podocarpus lambertii* (7.9%) and Poaceae (5.2%).

297 *Eugenia*-type, 'cf *Lithraea* a (Anacardiaceae)', Asteraceae (others total), *Lamanonia*, *I. theezans*,
298 *Calyptrothos*-type, *Vernonanthura*-type and *Clethra* pollen all had average abundances between
299 1% and 5%.

300 [Insert Figure 2 here]

301 Figure 2: Relative abundance (%) of selected tree species in the vegetation survey. Species included
302 in the cluster analysis and DCA are indicated with an asterisk.

303 [Insert Figure 3 here]

304 Figure 3: Relative Coverage Values (RCVs, %) of selected tree species in the vegetation survey. 'Ave.
305 (all)' = sum of RCVs/total number of plots; 'Ave. (present)' = sum of RCVs/number of plots in which
306 taxon was present.

307 [Insert Figure 4 here]

308 Figure 4: Relative abundance (%) of selected pollen types. Taxa included in the cluster analysis and
309 DCA are indicated with an asterisk

310 3.3. Plot differentiation

311 33 taxa from the vegetation survey and 15 pollen types met the 2% abundance threshold for
312 inclusion in the plot differentiation analysis (see section 2.4.2); these are identified with asterisks in
313 the summary diagrams (Figures 2 and 4).

314 [Insert Figure 5 here]

315 Figure 5: Cluster analyses (left; a, c) and DCAs (right; b, d) of plots by vegetation (top; a, b) and
316 pollen (bottom; c, d) taxa found in ≥ 2 plots at $\geq 2\%$ abundance. In the DCAs, convex hulls are
317 delimited for structural categories containing three or more plots. Plot types are identified by
318 colour and two-letter code (op: open; cl: closed; ri: riverine; di: disturbed; ro: rock; sl: slope; see

319 Table 1).

320 In the cluster analysis of vegetation data (Figure 5a), similarity scores range from ~0.25 to > 0.75.

321 Two plot groupings that are somewhat structurally consistent can be highlighted. One such cluster
322 contains three of the five open-understorey plots (P119, P122 and P123) with the lone slope plot
323 P118 (similarity score > 0.65); the other is the pairing of riverine plots P136 and P110 which, despite
324 having similarity scores < 0.6 relative to each other, are together highly dissimilar to all other plots.

325 In the DCA of vegetation data (Figure 5b), most variation is explained on axis 1 (eigenvalue 0.3781),
326 with values ranging from -196 (*Cyathea* sp.) to 365 (*Nectandra megapotamica*). The majority of
327 sample plots are clustered between 0 and 143 on axis 1, with the riverine plots P136 (202) and P110
328 (255) more distinguished. The values on axis 2 (eigenvalue 0.1268) range from -337 (*Sapium*
329 *glandulosum*) to 388 (*Myrsine umbellata*); sample plots are all clustered between 0 and 140 on this
330 axis.

331 In the pollen cluster analysis (Figure 5c), plots' similarity scores range from ~0.5 to > 0.85. Two small
332 clusters dominated by open understorey forest plots were found: P122 and P123 (both open;
333 similarity > 0.7), and P144 and P119 (both open) with the rocky forest plot P117 (similarity > 0.7).
334 These two groupings are quite dissimilar from each other, however, and the cluster that contains
335 them both also contains seven non-open plots, and excludes the final open plot (P142).

336 In the DCA of pollen spectra (Figure 5d) axis 1 (eigenvalue 0.2454) ranges from -72 (*Pteris*) to 305
337 (*Lamanonia*), with 15 of the plots clustered between 0 and 119; P151 is located at 184. The forest
338 plots are less well distinguished on axis 2 (eigenvalue 0.07194), sitting between values of 0 and 110
339 on an axis that ranges from -233 (Myrtaceae others total) and 426 (*Clethra* total).

340 3.4. Vegetation-pollen relationships.

341 We examined the vegetation-pollen relationships of 27 arboreal genera. Four of these

(*Myrceugenia*, *Nectandra*, *Ocotea* and *Piptocarpha*) were present in the vegetation but with no identifiable pollen produced or preserved ($p/v = 0$), and one (*Clethra*) was frequent in the pollen record without being recorded in the vegetation survey (giving it a mathematically infinite p/v value). Four other genera had higher average abundances for pollen than vegetation ($p/v > 1$), and 18 were more abundant in the vegetation than pollen ($p/v < 1$) – see Figure 6.

Of the taxa found in both pollen and vegetation datasets, the most over-represented genus is *Lamanonia*, followed by *Podocarpus*, *Myrsine* and *Araucaria*. In plot P151 *Lamanonia* was not recorded in the vegetation but its pollen made up 20.3% of the total; with this data point excluded from the average calculations, *Lamanonia*'s p/v value is 1.25. The most under-represented genera found in the pollen rain are *Cinnamodendron*, *Allophylus*, *Prunus*, *Styrax*, *Luehea* and *Zanthoxylum* ($p/v < 0.05$); *Casearia*, *Matayba*, *Pera* and *Lithraea* also have p/v scores < 0.1 . *Calypttranthes*, *Campomanesia*, *Eugenia*, *Sapium*, *Sebastiania* and *Xylosma* all have p/v values between 0.1 and 0.5. The pollen type 'cf *Lithraea* a (Anacardiaceae)' was three times more common than pollen that could be confidently identified to the *Lithraea* genus; assigning these grains to *Lithraea* changes the genus's p/v score from 0.09 to 0.39.

[Insert Figure 6 here]

Figure 6: p/v values of key genera of *Araucaria* forest in this study. *Clethra* has an infinitely high p/v value; faint sections in the bars of *Lamanonia* and *Lithraea* reflect alternate calculations of their p/v values (see section 3.4).

Between-plot variability in vegetation-pollen relationships is not well demonstrated by p/v values, as their calculation requires a taxon to be present in both vegetation and pollen records for a plot. The only taxon for which this occurs in all plots is *Eugenia*, whose plot-specific p/v scores varied between 0.04 (P141) and 2.26 (P122), with 14 plots having p/v values < 1 . This variability is more

365 evident with pollen-vegetation abundance differentials (Table 3 and Figure 7). *Myrsine*, *Araucaria*
 366 and *Podocarpus* had the most positive mean values (all above +5%); *Nectandra*, *Eugenia*, *Lithraea*,
 367 *Cinnamodendron*, *Styrax*, *Ocotea*, and *Matayba* had the most negative mean values (all below -5%).
 368 *Lithraea*'s mean differential changes to -5.52% if 'cf *Lithraea* (Anacardiaceae)' pollen is included in
 369 the calculation.

Genus	Mean	Min	Max	Range (Max-Min)
<i>Allophylus</i>	-4.13	-8.57	0.20	8.77
<i>Araucaria</i>	+9.36	-9.48	20.63	30.11
<i>Calyptranthes</i>	-3.44	-17.06	1.09	18.15
<i>Campomanesia</i>	-2.08	-5.71	0.81	6.52
<i>Casearia</i>	-4.17	-13.84	0.45	14.29
<i>Cinnamodendron</i>	-8.08	-24.14	0.44	24.57
<i>Clethra</i>	+1.76	0.37	6.68	6.31
<i>Eugenia</i>	-8.47	-25.92	7.17	33.09
<i>Ilex</i>	-1.07	-8.73	2.93	11.66
<i>Lamanonia</i>	+2.69	-1.16	20.32	21.49
<i>Lithraea</i>	-8.25	-32.56	0.26	32.82
<i>Luehea</i>	-4.89	-8.89	0.18	9.07
<i>Matayba</i>	-6.29	-34.14	0.36	34.50
<i>Myrceugenia</i>	-4.61	-9.09	-1.75	7.34

<i>Myrsine</i>	+13.88	-3.05	55.75	58.80
<i>Nectandra</i>	-11.73	-28.57	-2.17	26.40
<i>Ocotea</i>	-7.25	-16.36	-1.75	14.61
<i>Pera</i>	-4.83	-21.04	1.21	22.25
<i>Piptocarpha</i>	-3.02	-4.29	-1.75	2.53
<i>Podocarpus</i>	+5.78	-5.73	19.24	24.97
<i>Prunus</i>	-3.21	-6.82	0.23	7.04
<i>Sapium</i>	-1.27	-5.05	0.44	5.49
<i>Sebastiania</i>	-4.35	-25.85	0.55	26.40
<i>Styrax</i>	-7.52	-22.86	0.45	23.31
<i>Vernonanthura</i>	-0.38	-4.89	1.83	6.72
<i>Xylosma</i>	-2.16	-7.02	1.49	8.51
<i>Zanthoxylum</i>	-4.76	-12.73	0.20	12.93

370 Table 3: Pollen-vegetation abundance differential values (%) for key genera of *Araucaria* forest in
371 this study.

372 *Myrsine* had the most extreme positive values: +55.75% (P114, slope), +41.25% (P157, disturbed),
373 +30.70% (P141, closed understorey), +26.38% (P142, open understorey) and +20.48% (P159,
374 disturbed). Values above +20% also occurred twice for *Araucaria* (+20.63% in P151, riverine, and
375 +20.30% in P159, disturbed) and once for *Lamanonia* (+20.32% in P151, riverine). Similarly negative
376 values (below -20%) were more evenly spread among taxa, with two cases in *Styrax* (-22.86% in
377 P122, open understorey, and -20.00% in P141, closed understorey) and one each in *Matayba* (-

378 34.14% in P136, riverine), *Lithraea* (-32.56% in P114, slope), *Nectandra* (-28.57% in P110, riverine),
379 *Eugenia* (-25.92% in P156, disturbed), *Sebastiania* (-25.85% in P151, riverine), *Cinnamodendron* (-
380 24.14% in P159, disturbed), and *Pera* (-21.04% in P119, open understorey).

381 [Insert Figure 7 here]

382 Figure 7: Pollen-vegetation abundance differentials (square-root transformed) for selected genera.
383 Positive values denote plots where a taxon's pollen was more abundant than its vegetation;
384 negative values reflect vegetation being more abundant than pollen. Blank cells denote plots where
385 a taxon was absent from both pollen and vegetation data.

386 The genus with the greatest difference between maximum and minimum abundance differentials is
387 *Myrsine*, with wide variation (> 30%) also found in *Matayba*, *Eugenia*, *Lithraea* and *Araucaria*;
388 *Nectandra*, *Sebastiania*, *Podocarpus*, *Cinnamodendron*, *Styrax* and *Lamanonia* had differences
389 between 20 and 30%. Most genera had both positive and negative abundance differentials, with
390 the exceptions being *Clethra* (all positive) and *Myrceugenia*, *Nectandra*, *Ocotea* and *Piptocarpha* (all
391 negative). However, few taxa (*Araucaria*, *Eugenia*, *Ilex*, *Myrsine*, *Podocarpus*) had differentials > 2%
392 in both positive and negative directions.

393 4. Discussion

394 4.1. Detecting structural differences in *Araucaria* forest

395 The vegetation of the studied forest area is characteristic of *Araucaria* forest, containing 19 of its 30
396 most important tree species (Schorn et al., 2012). Three of the predominant species in this study (*A.*
397 *angustifolia*, *O. pulchella*, *P. lambertii*), as well as others found at lower levels (*Matayba*
398 *elaeagnoides*, *Lamanonia ternata*), are key indicator species (Gonçalves and Souza, 2014). *Lithraea*
399 *brasiliensis*, the most common, abundant and dominant species in our sample plots, is the fifth

most important tree across Santa Catarina's *Araucaria* forest, especially significant below 1200 m a.s.l (Meyer et al., 2013; Schorn et al., 2012).

The floristic composition of the plots is somewhat variable, with no pair of plots having a similarity score above 0.8 (Figure 5a), but this variability in composition does not generally reflect the structural differences between the plots. This can be seen in the cluster analysis (Figure 5a): three of the open understorey plots (P119, P122, P123) are clustered together along with the slope plot P118, but the remaining two open plots (P142 and P144) are far removed; the cluster that would include all five open plots would have 14 members. Similarly, the disturbed plots P157 and P159 are closely paired, but the cluster that would also include the third disturbed plot (P156) would encompass nine plots in total. The riverine plots P110 and P136 are distinct from the rest of the plots but cannot be grouped with the third member of the riverine group (P151) without including all sampled plots in the cluster.

This pattern is further shown in the DCA (Figure 5b), of which only axis 1 has an eigenvalue > 0.3 (0.3781), which would be suggestive of ecological relevance (Peterson and Gale, 1991; Rezende et al., 2015). Here again P110 and P136 are the only plots that are clearly distinguished by their vegetation. Their floristic difference to the other studied plots may partly result from their location at the northern end of the studied area, > 1.5 km from the other plots. Their regeneration stage could also be an explanatory factor: these plots have the highest RCVs for *Matayba elaeagnoides*, as well as the highest recorded abundances of the pioneer *Nectandra megapotamica*, two of the three highest abundances of *Allophylus edulis*, and the presence of *Sebastiania commersoniana*, all of which are important taxa in regenerating *Araucaria* forest patches (Meyer et al., 2013; Schorn et al., 2012).

The most common and abundant tree pollen in our study sites came from *Araucaria angustifolia*,

423 Myrtaceae (mainly *Eugenia*), Anacardiaceae cf *Lithraea*, *Myrsine* and *Podocarpus*, with *Lithraea*,
424 *Vernonanthura*-type, *Sebastiania*, *Ilex* and *Clethra* found in the majority of plots at generally low
425 abundances (Figure 4). Pteridophyte spores were both common and abundant, and Poaceae was
426 the most prevalent herbaceous pollen taxon, with Asteraceae and Araceae pollen also common but
427 less abundant. The sum of all herbaceous pollen varies from 5-20%, averaging 10% – similar to the
428 5-15% found by Garcia et al. (2004) in a study of *Araucaria* forest peat deposits in Sao Paulo state,
429 but lower than the 24-60% found by Jeske-Pieruschka et al. (2010) in a forest patch surrounded by
430 campos grassland. The herbaceous pollen counts in the latter were likely increased by influx from
431 the grasslands surrounding the forest, and as a result of disturbance from grazing in the studied
432 forest patch (Jeske-Pieruschka et al., 2010).

433 The plots are less well differentiated by pollen than by their vegetation, as shown by their generally
434 higher cluster similarity scores (Figure 5c). As with the vegetation there are some potential pairings
435 of open understorey plots (P122 and P123; P119 and P144 with the rocky plot P117), but these are
436 quite distinct from each other – the cluster that would encompass all five open plots would only
437 exclude the riverine plot P151. No potential groupings in the pollen cluster analysis mirror clusters
438 of the plots' vegetation (Figure 5a,c), and there are no consistent groupings of structurally similar
439 plots. This is also seen in the DCA (Figure 5d), where the eigenvalue for axis 1 (0.2454) may indicate
440 the separation is not ecologically relevant (Peterson and Gale, 1991; Rezende et al., 2015). Here, as
441 in the cluster analysis, the only plot that is clearly distinguished is P151, due in large part to its
442 exceptionally high abundance of *Lamanonia* pollen.

443 These results show that the notable structural differences between the different forest plot
444 categories are not echoed in the composition of their woody taxa. Since there are no consistent
445 floristic differences between plot types, it is unsurprising that these structural differences cannot be

446 detected by their pollen signals – especially as pollen’s sensitivity is affected by relatively coarse
447 taxonomic resolution and differential pollen production and preservation between taxa.

448 4.2. Vegetation-pollen relationships

449 The pollen records of the studied forest plots did not consistently reflect large variations in a
450 taxon’s abundance in the vegetation (Figure 6 & 7). *L. brasiliensis*, for example, makes up 2.9% of
451 the vegetation in P110 and 34.1% in P114 (Figure 2), but *Lithraea* pollen is equally abundant in each
452 plot (1.6%) (Figure 4); its abundance in the vegetation is intermediate in P117 (10.9%) and P118
453 (12.3%) but its pollen makes up just 0.9% and 1.0% of those plot totals, respectively. This variability
454 in vegetation-pollen relationships can also be illustrated with *Myrsine*: its pollen made up more
455 than half of all counted grains (55.8%) in P114, despite no individuals from the genus being
456 recorded in the vegetation survey; it was also significantly over-represented in P157 (pollen
457 abundance 43.4%, vegetation abundance 2.2%), but proportionately represented in P119 (17.3% of
458 the pollen; 17.0% of the vegetation) and under-represented in P118 (5.7% of the pollen, 8.8% of the
459 vegetation).

460 However, general patterns of over- or under-representation of each taxon in the pollen record
461 could be determined – only *Araucaria*, *Eugenia*, *Ilex*, *Myrsine* and *Podocarpus* had both positive and
462 negative abundance differentials above 2% (Table 3). The majority (22 of 27) of the arboreal genera
463 examined were under-represented in the pollen record (p/v values <1; Figure 6), with four
464 (*Myrceugenia*, *Nectandra*, *Ocotea* and *Piptocarpha*) being palynologically silent. These under-
465 represented taxa include many of the most abundant tree species in the plots, as well as some of
466 the most ecologically important taxa in *Araucaria* forest (*Lithraea*, *Matayba*, *Cinnamodendron*,
467 *Ocotea*, *Prunus* and *Nectandra*) (Sevegnani et al., 2013). The highest p/v value among these key
468 genera is *Lithraea*’s 0.09 (i.e. less than one tenth as abundant in the pollen record as in the

469 vegetation), and no pollen which could be reliably identified as *Ocotea* or *Nectandra* was counted
470 at all.

471 Under-representation in the pollen record is common in the Neotropics (Bush, 1995; Bush and
472 Rivera, 1998; Gosling et al., 2009; Haselhorst et al., 2013). This is partly due to the relative rarity of
473 anemophilous plants – generally the most over-represented taxa – in tropical forests, although
474 entomophilous plants are less under-represented in tropical pollen spectra than in temperate
475 regions (Bush, 1995; Viera et al., 2012; Whitehead, 1983). Our results fit this general pattern: of the
476 under-represented taxa in our study, *Nectandra* and *Ocotea* are pollinated by thrips (Thysanoptera)
477 (Danieli-Silva and Varassin, 2013; Souza and Moscheta, 2000), and *Lithraea*, *Ilex*, *Matayba*, *Prunus*,
478 *Eugenia* and *Cinnamodendron* depend on bees and other small insects (de Deus et al., 2014;
479 Hermes and Köhler, 2006; Imperatriz-Fonseca et al., 1989; Montalva et al., 2011; Tomlinson, 1974;
480 Wilms et al., 1997). Additionally, *Ocotea* pollen is fragile, with a thin exine that means it preserves
481 poorly (Behling, 1993). By contrast, the over-represented taxa *Araucaria* (Bittencourt and Sebbenn,
482 2008, 2007) and *Podocarpus* (Midgley, 1989; Negash, 2003; Wilson and Owens, 1999) are
483 anemophilous, as are South American members of *Myrsine* (Albuquerque et al., 2013; Otegui and
484 Cocucci, 1999), and therefore produce abundant pollen.

485 *Lamanonia* was over-represented in the pollen record (p/v 5.44; 1.25 when excluding P151) despite
486 being insect-pollinated (de Deus et al., 2014; Hermes and Köhler, 2006; Wilms et al., 1997). Its
487 relative abundance of pollen in this study may be explained by its mass-flowering habit (Hermes
488 and Köhler, 2006), so it is possible that *Lamanonia*'s pollen-vegetation relationship varies
489 significantly year-on-year. *Clethra* is another entomophilous taxon (Freitas and Sazima, 2006) with
490 over-represented pollen in this study: its pollen was found in 10 of the 16 plots at an average
491 abundance of 1.10% (0.37-4.58%), but no individuals were recorded in the vegetation (Figure 2)

despite *C. scabra* being the third most important tree species in *Araucaria* forest (Schorn et al., 2012). Given *Clethra* pollen is not adapted for long-distance travel, it is unlikely to have originated outside the plots; potentially, individuals from this genus were present but too small to be included in the survey (i.e. d.b.h. < 10 cm), making it difficult to make a true assessment of the taxon's pollen-vegetation relationship.

4.3. Implications for palaeoecology

Our findings have important implications for the interpretation of fossil pollen records. Many of the key taxa used to reconstruct past *Araucaria* forest dynamics were identified palynologically in this study, but two were largely missing. Pollen from *Mimosa scabrella* and *Weinmannia* is commonly cited as evidence of *Araucaria* forest (Behling, 1997a, 1995; Behling et al., 2004; Leonhardt and Lorscheitter, 2010) but *M. scabrella* pollen was not abundant (present in 10 of the 16 plots, maximum abundance 0.89%) and *Weinmannia* pollen was absent; neither species was recorded in the vegetation survey. *M. scabrella* is a species which is especially encouraged by anthropogenic forest disturbance, so its absence here may be connected to the relatively long time since the last anthropogenic disturbance of our studied plots. Moreover, both taxa (especially *Weinmannia*) are much more floristically important at elevations above 1,200 m a.s.l. (Sevegnani et al., 2013) so their absence here is unsurprising, and highlights the biases of existing palaeoecological studies towards high-altitude regions along the eastern edge of the highland plateau (Behling, 2007, 1997a, 1995; Behling et al., 2004; Jeske-Pieruschka et al., 2013; Leonhardt and Lorscheitter, 2010; Scherer and Lorscheitter, 2014).

The geographical bias of existing studies can further be seen in the classifications proposed by Rodrigues et al. (2016b) for distinguishing woodland types in southern Brazil based on pollen spectra. Although our sites (721-953 m a.s.l.) have vegetation which is most similar to that in the

515 'low Poaceae subtropical forest' (LPSF) category (Behling et al., 2004; Behling and de Oliveira, 2018;
516 Jeske-Pieruschka et al., 2013; Jeske-Pieruschka and Behling, 2012), they are excluded from this
517 group due to their absence of *Weinmannia* pollen. Instead, our plots would likely be included in the
518 disparate 'low Poaceae subtropical-tropical' (LPST) group, along with two sites in Santa Catarina's
519 coastal tropical forest (9-10 m a.s.l.) (Behling, 1997b, 1995), one in northern Argentina at the
520 westernmost limit of *Araucaria* forest taxa (604 m a.s.l.) (Gessert et al., 2011), and another at low
521 altitude in south-central Rio Grande do Sul (176 m a.s.l.) (Santa Mônica, unpublished, cited in
522 Rodrigues et al., 2016b) . The significant altitudinal, geographical and floristic differences between
523 these sites suggests LPST may not be a true grouping, and that the classification of southern Brazil's
524 forests by their pollen spectra will improve as more westerly plateau sites, with different
525 assemblages of *Araucaria* forest, are studied.

526 Most of the taxa widely used in palaeoecological studies were here found to be over-represented in
527 the pollen record (*Araucaria*, *Myrsine*, *Podocarpus*, *Lamanonia*) or under-represented but very
528 abundant in the vegetation (Myrtaceae, such as *Eugenia* and *Calypttranthes* which are rarely
529 distinguished to genus in fossil pollen). Fossil *Lithraea* pollen is quite often identified (Behling, 2007;
530 Behling et al., 2016, 2005; Gessert et al., 2011) or grouped with *Schinus* (Behling, 1997b, 1997a;
531 Behling et al., 2004; Jeske-Pieruschka et al., 2010) but rarely treated as important (but see Behling,
532 1997a; Gessert et al., 2011); given the floristic importance of *L. brasiliensis* (Meyer et al., 2013;
533 Schorn et al., 2012) and its under-representation in the pollen rain (p/v 0.09-0.39, pollen 8.25% less
534 abundant than vegetation on average), the presence and dynamics of *Lithraea* pollen may warrant
535 more detailed examination in future studies. *Lithraea* is one of several key *Araucaria* forest taxa we
536 have shown to be either significantly under-represented (*Matayba*, *Cinnamodendron*, *Prunus*) or
537 absent (*Ocotea*, *Nectandra*) in the pollen record, even when quite abundant in the vegetation. This
538 suggests that some significant changes in past forest composition may not be reflected in fossil

539 pollen, highlighting a potentially important limitation of pollen as a palaeoecological proxy for
540 determining this ecosystem's responses to past millennial-scale human land use and climate
541 changes.

542 The importance of fossil pollen's limited sensitivity to compositional changes in *Araucaria* forest is
543 reinforced when considering that structural differences between plots were not clearly reflected in
544 their pollen signals. Although the plots had notably different structural characteristics to one
545 another, this was only weakly echoed in the floristic composition of their arboreal component, and
546 even less so in their pollen spectra, which were more similar to each other than the vegetation. This
547 implies that pollen records may not clearly detect structural changes in *Araucaria* forest, especially
548 when these are not accompanied by significant or long-term alterations in floristic composition.
549 Further investigations of the sensitivity of pollen rain to larger-scale changes in composition and
550 structure of *Araucaria* forest might involve integration with the state-wide plot network of the
551 Santa Catarina forest floristic inventory (Vibrans et al., 2010) or palynological examination of forest
552 fragments with relatively intense present human land use, such as caívas, faxinais or ervais (Fichino
553 et al., 2017; Mello and Peroni, 2015; Reis et al., 2018). Such research would aid the interpretation
554 of fossil pollen records which do show significant changes with climatic changes and human land
555 use (e.g. Behling et al., 2004), permitting shifts in pollen composition to be more accurately related
556 to changes in the vegetation.

557 The difficulty of detecting structural and compositional changes in *Araucaria* forest with pollen
558 records has implications for the reconstruction of past human impacts on the forest. Before
559 European arrival, the indigenous southern Jê people lived in semi-subterranean dwellings in the
560 forest, practising swidden cultivation and exploiting *Araucaria angustifolia* nuts and Myrtaceae
561 fruits (Corteletti, 2012; Corteletti et al., 2015; dos Reis et al., 2014; Iriarte and Behling, 2007). It has

562 been suggested that their lifestyles helped to spread *Araucaria* forest more quickly than natural
563 drivers, such as climate change, and even beyond the forest's natural limits (Bitencourt and
564 Krauspenhar, 2006; Iriarte and Behling, 2007; Lauterjung et al., 2018; Noelli, 2000; Robinson et al.,
565 2018), but our findings raise the question of whether, and how, more subtle impacts within the
566 forest – especially structural and compositional changes – would be represented in fossil pollen
567 sedimentary archives. There is significant ongoing debate over the extent to which pre-Columbian
568 peoples altered the floristic composition of Amazonian forests (Barlow et al., 2012; Levis et al.,
569 2018, 2017; McMichael et al., 2017; Shepard and Ramirez, 2011) and the limitations of fossil pollen
570 analysis in identifying such impacts are well known (e.g. Carson et al., 2016), with recent
571 simulations showing that pollen sites in tropical forest-grassland mosaics (like those between
572 *Araucaria* forest and campos grasslands) may be insensitive to large changes in forest cover
573 (Whitney et al., 2019). The forest plots in our study have been free of human disturbance for
574 several decades, so their present condition is unlikely to mirror the full range of southern Jê land
575 use. Nevertheless, our findings highlight potential limitations of pollen analysis as a tool for
576 investigating and reconstructing the role of the southern Jê in shaping the *Araucaria* forest's
577 floristic composition and structure – the absence of evidence of human impacts may not equate to
578 evidence of their absence.

579 5. Conclusions

580 In this study, the pollen rain from 16 structurally different *Araucaria* forest plots has been
581 characterised and compared with floristic inventory data, and 27 tree genera have had their
582 vegetation-pollen relationships quantified.

583 Our results show that the structurally different plots do not exhibit consistent differences in the
584 floristic composition of their tree taxa. Correspondingly, their structural differences cannot be

585 clearly detected in their pollen rain, with pollen spectra more similar than vegetation between
586 plots.

587 On average, the most abundant tree species in the plots are *Lithraea brasiliensis*, *Cinnamodendron*
588 *dinisii*, *Araucaria angustifolia*, *Eugenia* sp., *Ocotea pulchella* and *Calyptranthes* cf. *concinna*,
589 representing the typical composition of *Araucaria* forest at this altitude. As well as these, *Styrax*
590 *leprosus* has a high average relative coverage value (RCV) across all plots, and species including
591 *Matayba elaeagnoides*, *Nectandra megapotamica*, *Vernonanthura discolor*, *Prunus myrtifolia*,
592 *Piptocarpha angustifolia*, *Myrceugenia* sp. and *Myrsine umbellata* have high RCVs in the (relatively
593 few) plots in which they occur. The modern pollen rain of *Araucaria* forest in our study area (720-
594 920m a.s.l.) is characterised by *Myrsine* (19.5%), *A. angustifolia* (16.6%), Pteridophyta (13.2%),
595 *Podocarpus lambertii* (7.9%) and Poaceae (5.2%), along with *Eugenia*-type, Anacardiaceae cf.
596 *Lithraea*, Asteraceae (others), *Lamanonia*, *Ilex theezans*, *Calyptranthes*-type, *Vernonanthura*-type
597 and *Clethra* at lower abundance.

598 Vegetation-pollen relationships vary between plots but general patterns of over- or under-
599 representation could be identified for the majority of taxa. *Araucaria*, *Lamanonia*, *Podocarpus*, and
600 *Myrsine* are over-represented in the pollen record (average $p/v > 1$), and *Clethra* pollen was
601 frequently identified but the genus is absent from the vegetation survey. All of the other 22 genera
602 analysed are under-represented, with *Myrceugenia*, *Nectandra*, *Ocotea* and *Piptocarpha*
603 palynologically silent.

604 The palynological under-representation of ecologically key *Araucaria* forest taxa and the lack of
605 clear differentiation of structural differences in the forest's pollen spectra have important
606 implications for the interpretation of fossil pollen records, highlighting the challenges of resolving
607 past human impacts and ecological changes within *Araucaria* forest using this proxy. This

608 quantitative analysis of vegetation-pollen relationships for key *Araucaria* forest taxa is a key tool for
609 interpreting fossil pollen records and understanding past vegetation dynamics on the highlands of
610 southern Brazil.

611 Author contributions

612 MLC and FEM conceived the study; MLC collected moss polsters; MLC and LAS collected vegetation
613 survey data; MLC performed pollen analysis and counting; MLC and OJW analysed the data and
614 interpreted the results; OJW, MLC and FEM drafted the manuscript; all authors commented on the
615 manuscript before submission.

616 Acknowledgements

617 This study was funded by an AHRC-FAPESP research grant entitled ‘Je Landscapes of southern
618 Brazil: Ecology, History and Power in a transitional landscape during the Late Holocene’
619 (AH/K004212/1) to JI and FEM, with MLC as appointed PDRA with FEM; OJW was funded by a
620 University of Reading Graduate Teaching Assistant PhD studentship. The authors thank the Florestal
621 Gateados Company for permission to conduct this research on their property; Museu Botânico
622 Municipal of Curitiba for providing key herbarium samples for modern pollen references; Alyne
623 Ruggiero and Alvaro Cortes for fieldwork assistance; Professor Hermann Behling for assistance with
624 pollen identification; Dr John Carson for constructive comments in the early stages of the study;
625 and Dr Mark Robinson for early editing support.

626 References

627 Albuquerque, A.A.E., de Lima, H.A., Gonçalves-Esteves, V., Benevides, C.R., Rodarte, A.T.A., 2013.
628 *Myrsine parvifolia* (Primulaceae) in sandy coastal plains marginal to Atlantic rainforest: a case
629 of pollination by wind or by both wind and insects? *Brazilian J. Bot.* 36, 65–73.

630 <https://doi.org/10.1007/s40415-013-0007-4>

631 Alvares, C.A., Stape, J.L., Sentelhas, P.C., De Moraes Gonçalves, J.L., Sparovek, G., 2013. Köppen's
632 climate classification map for Brazil. *Meteorol. Zeitschrift* 22, 711–728.
633 <https://doi.org/10.1127/0941-2948/2013/0507>

634 Atanassova, J.R., 2007. Pollen deposition in moss polsters and pollen traps in the Central Stara
635 Planina Mts (2002–2005). *Phytol. Balc.* 13, 223–228.

636 Barlow, J., Gardner, T.A., Lees, A.C., Parry, L., Peres, C.A., 2012. How pristine are tropical forests? An
637 ecological perspective on the pre-Columbian human footprint in Amazonia and implications for
638 contemporary conservation. *Biol. Conserv.* 151, 45–49.
639 <https://doi.org/10.1016/j.biocon.2011.10.013>

640 Behling, H., 2007. Late Quaternary vegetation, fire and climate dynamics of Serra do Aracatuba in
641 the Atlantic coastal mountains of Parana State, southern Brazil. *Veg. Hist. Archaeobot.* 16, 77–
642 85. <https://doi.org/10.1007/s00334-006-0078-2>

643 Behling, H., 2002. South and southeast Brazilian grasslands during Late Quaternary times: A
644 synthesis. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 177, 19–27.
645 [https://doi.org/10.1016/S0031-0182\(01\)00349-2](https://doi.org/10.1016/S0031-0182(01)00349-2)

646 Behling, H., 1997a. Late Quaternary vegetation, climate and fire history of the Araucaria forest and
647 campos region from Serra Campos Gerais, Paraná State (South Brazil). *Rev. Palaeobot. Palynol.*
648 97, 109–121. [https://doi.org/10.1016/S0034-6667\(96\)00065-6](https://doi.org/10.1016/S0034-6667(96)00065-6)

649 Behling, H., 1997b. Late Quaternary vegetation, climate and fire history from the tropical mountain
650 region of Morro de Itapeva, SE Brazil. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 129, 407–422.
651 [https://doi.org/10.1016/S0031-0182\(97\)88177-1](https://doi.org/10.1016/S0031-0182(97)88177-1)

- 652 Behling, H., 1995. Investigations into the late Pleistocene and Holocene history of vegetation and
653 climate in Santa Catarina (S Brazil). *Veg. Hist. Archaeobot.* 4, 127–152.
654 <https://doi.org/10.1007/BF00203932>
- 655 Behling, H., 1993. Untersuchungen zur Spatpleistozänen und Holozänen Vegetations- und
656 Klimageschichte der Tropischen Küstenwälder und der Araukarienwälder in Santa Catarina
657 (Sudbrasilien). Diss. Bot. J. Cramer, Stuttgart, Germany.
- 658 Behling, H., Bauermann, S.G., Neves, P.C.P. das, 2001. Holocene environmental changes in the Sao
659 Francisco de Paula region, southern Brazil. *J. South Am. Earth Sci.* 14, 631–639.
660 [https://doi.org/10.1016/S0895-9811\(01\)00040-2](https://doi.org/10.1016/S0895-9811(01)00040-2)
- 661 Behling, H., de Oliveira, M.A.T., 2018. Evidence of a late glacial warming event and early Holocene
662 cooling in the southern Brazilian coastal highlands. *Quat. Res.* 89, 90–102.
663 <https://doi.org/10.1017/qua.2017.87>
- 664 Behling, H., Pillar, V.D., Bauermann, S.G., 2005. Late Quaternary grassland (Campos), gallery forest,
665 fire and climate dynamics, studied by pollen, charcoal and multivariate analysis of the São
666 Francisco de Assis core in western Rio Grande do Sul (southern Brazil). *Rev. Palaeobot. Palynol.*
667 133, 235–248. <https://doi.org/10.1016/j.revpalbo.2004.10.004>
- 668 Behling, H., Pillar, V.D., Orlóci, L., Bauermann, S.G., 2004. Late Quaternary Araucaria forest,
669 grassland (Campos), fire and climate dynamics, studied by high-resolution pollen, charcoal and
670 multivariate analysis of the Cambará do Sul core in southern Brazil. *Palaeogeogr.*
671 *Palaeoclimatol. Palaeoecol.* 203, 277–297. [https://doi.org/10.1016/S0031-0182\(03\)00687-4](https://doi.org/10.1016/S0031-0182(03)00687-4)
- 672 Behling, H., Verissimo, N., Bauermann, S., Bordignon, S., Evaldt, A., 2016. Late Holocene Vegetation
673 History and Early Evidence of *Araucaria angustifolia* in Caçapava do Sul in the Lowland Region

674 of Rio Grande do Sul State, Southern Brazil. Brazilian Arch. Biol. Technol. 59.
 675 <https://doi.org/10.1590/1678-4324-2016150264>

676 Bergamin, R.S., 2017. Variação espacial e temporal de espécies arbóreas em florestas da Mata
 677 Atlântica: perspectivas sobre o efeito de mudanças climáticas e planejamento de conservação.
 678 Universidade Federal do Rio Grande do Sul.

679 Bernal, J.P., Cruz, F.W., Stríkis, N.M., Wang, X., Deininger, M., Catunda, M.C.A., Ortega-Obregón, C.,
 680 Cheng, H., Edwards, R.L., Auler, A.S., 2016. High-resolution Holocene South American monsoon
 681 history recorded by a speleothem from Botuverá Cave, Brazil. Earth Planet. Sci. Lett. 450, 186–
 682 196. <https://doi.org/10.1016/j.epsl.2016.06.008>

683 Bittencourt, A.L.V., Krauspenhar, P.M., 2006. Possible prehistoric anthropogenic effect on *Araucaria*
 684 *angustifolia* (Bert.) O. Kuntze expansion during the late Holocene. Rev. Bras. Paleontol. 9, 109–
 685 116.

686 Bittencourt, J.V.M., Sebbenn, A.M., 2008. Pollen movement within a continuous forest of wind-
 687 pollinated *Araucaria angustifolia*, inferred from paternity and TwoGener analysis. Conserv.
 688 Genet. 9, 855–868. <https://doi.org/10.1007/s10592-007-9411-2>

689 Bittencourt, J.V.M., Sebbenn, A.M., 2007. Patterns of pollen and seed dispersal in a small,
 690 fragmented population of the wind-pollinated tree *Araucaria angustifolia* in southern Brazil.
 691 Heredity (Edinb). 99, 580–591. <https://doi.org/10.1038/sj.hdy.6801019>

692 Burn, M.J., Mayle, F.E., Killeen, T.J., 2010. Pollen-based differentiation of Amazonian rainforest
 693 communities and implications for lowland palaeoecology in tropical South America.
 694 Palaeogeogr. Palaeoclimatol. Palaeoecol. 295, 1–18.
 695 <https://doi.org/10.1016/j.palaeo.2010.05.009>

696 Bush, M.B., 1995. Neotropical Plant Reproductive Strategies and Fossil Pollen Representation. *Am.*
697 *Nat.* 145, 594–609. <https://doi.org/10.1086/285757>

698 Bush, M.B., Rivera, R., 1998. Pollen Dispersal and Representation in a Neotropical Rain Forest. *Glob.*
699 *Ecol. Biogeogr. Lett.* 7, 379–392. <https://doi.org/10.2307/2997685>

700 Bush, M.B., Weng, C., 2007. Introducing a new (freeware) tool for palynology. *J. Biogeogr.* 34, 377–
701 380. <https://doi.org/10.1111/j.1365-2699.2006.01645.x>

702 Carson, J.F., Mayle, F.E., Whitney, B.S., Iriarte, J., Soto, J.D., 2016. Pre-Columbian ring ditch
703 construction and land use on a “chocolate forest island” in the Bolivian Amazon. *J. Quat. Sci.*
704 31, 337–347. <https://doi.org/10.1002/jqs.2835>

705 Carvalho, M.M.X. de, Nodari, E.S., 2010. As fases da exploração madeireira na floresta com
706 Araucária e os progressivos avanços da indústria madeireira sobre as florestas primárias (1870-1970), in: Simpósio Internacional de História Ambiental e Migrações. UFSC,
707 Florianópolis, pp. 707–726.

709 Caseldine, C.J., 1989. Pollen assemblage–plant community relationships on the Storbreen glacier
710 foreland, Jotunheimen Mountains, southern Norway. *New Phytol.* 111, 105–118.
711 <https://doi.org/10.1111/j.1469-8137.1989.tb04224.x>

712 Chou, S.C., Lyra, A., Mourão, C., Dereczynski, C., Pilotto, I., Gomes, J., Bustamante, J., Tavares, P.,
713 Silva, A., Rodrigues, D., Campos, D., Chagas, D., Sueiro, G., Siqueira, G., Marengo, J., 2014.
714 Assessment of Climate Change over South America under RCP 4.5 and 8.5 Downscaling
715 Scenarios. *Am. J. Clim. Chang.* 03, 512–527. <https://doi.org/10.4236/ajcc.2014.35043>

716 Colinvaux, P., Oliveira, P.E. de, Moreno Patiño, J.E., 1999. Amazon pollen manual and atlas.
717 Harwood Academic Publishers.

718 Colombo, A.F., Joly, C.A., 2010. Brazilian Atlantic Forest lato sensu: the most ancient Brazilian
719 forest, and a biodiversity hotspot, is highly threatened by climate change. *Brazilian J. Biol.* 70,
720 697–708. <https://doi.org/10.1590/S1519-69842010000400002>

721 Correa-Metrio, A., Dechnik, Y., Lozano-García, S., Caballero, M., 2014. Detrended correspondence
722 analysis: A useful tool to quantify ecological changes from fossil data sets. *Boletín la Soc.*
723 *Geológica Mex.* 66, 135–143.

724 Corteletti, R., 2012. Projeto Arqueológico Alto Canoas-Paraca. Um estudo da presença Jê no
725 planalto catarinense. Universidade de São Paulo.

726 Corteletti, R., Dickau, R., DeBlasis, P., Iriarte, J., 2015. Revisiting the economy and mobility of
727 southern proto-Jê (Taquara-Itararé) groups in the southern Brazilian highlands: Starch grain
728 and phytoliths analyses from the Bonin site, Urubici, Brazil. *J. Archaeol. Sci.* 58, 46–61.
729 <https://doi.org/10.1016/j.jas.2015.03.017>

730 Danieli-Silva, A., Varassin, I.G., 2013. Breeding system and thrips (Thysanoptera) pollination in the
731 endangered tree *Ocotea porosa* (Lauraceae): Implications for conservation. *Plant Species Biol.*
732 28, 31–40. <https://doi.org/10.1111/j.1442-1984.2011.00354.x>

733 Davis, M.B., 1963. On the Theory of Pollen Analysis. *Am. J. Sci.* 261, 897–912.

734 de Deus, F.F., do Vale, V.S., Schiavini, I., Oliveira, P.E., 2014. Diversity of reproductive ecological
735 groups in semideciduous seasonal forests. *Biosci. J.* 30, 1885–1902.

736 De Souza, J.G., Corteletti, R., Robinson, M., Iriarte, J., 2016a. The genesis of monuments: Resisting
737 outsiders in the contested landscapes of southern Brazil. *J. Anthropol. Archaeol.* 41, 196–212.
738 <https://doi.org/10.1016/j.jaa.2016.01.003>

739 De Souza, J.G., Robinson, M., Corteletti, R., C??rdenas, M.L., Wolf, S., Iriarte, J., Mayle, F., DeBlasis,

740 P., 2016b. Understanding the chronology and occupation dynamics of oversized pit houses in
 741 the southern Brazilian highlands. PLoS One 11, 1–24.
 742 <https://doi.org/10.1371/journal.pone.0158127>

743 dos Reis, M.S., Ladio, A., Peroni, N., 2014. Landscapes with Araucaria in South America: Evidence for
 744 a cultural dimension. Ecol. Soc. 19, 43. <https://doi.org/10.5751/ES-06163-190243>

745 Duarte, L.D.S., Bergamin, R.S., Marcilio-Silva, V., Seger, G.D.D.S., Marques, M.C.M., 2014.
 746 Phylobetadiversity among forest types in the Brazilian Atlantic Forest complex. PLoS One 9.
 747 <https://doi.org/10.1371/journal.pone.0105043>

748 Faegri, K., Iversen, J., 1989. Textbook of Pollen Analysis, 4th ed, Textbook of pollen analysis. John
 749 Wiley & Sons Ltd., Chichester, UK.

750 Fernandes, R.C., Piovezana, L., 2015. The Kaingang perspectives on land and environmental rights in
 751 the south of Brazil. Ambient. Soc. 18, 111–128. [https://doi.org/10.1590/1809-](https://doi.org/10.1590/1809-4422ASOCEx07V1822015en)
 752 [4422ASOCEx07V1822015en](https://doi.org/10.1590/1809-4422ASOCEx07V1822015en)

753 Fichino, B.S., Pivello, V.R., Santos, R.F., 2017. Trade-offs among ecosystem services under different
 754 pinion harvesting intensities in Brazilian Araucaria Forests. Int. J. Biodivers. Sci. Ecosyst. Serv.
 755 Manag. 13, 139–149. <https://doi.org/10.1080/21513732.2016.1275811>

756 Forest, F., Moat, J., Baloch, E., Brummitt, N.A., Bachman, S.P., Ickert-Bond, S., Hollingsworth, P.M.,
 757 Liston, A., Little, D.P., Mathews, S., Rai, H., Rydin, C., Stevenson, D.W., Thomas, P., Buerki, S.,
 758 2018. Gymnosperms on the EDGE. Sci. Rep. 8, 6053. [https://doi.org/10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-24365-4)
 759 [24365-4](https://doi.org/10.1038/s41598-018-24365-4)

760 Formento, S., Schorn, L.A., Alaor, R., Ramos, B., 2004. Dinâmica Estrutural Arbórea De Uma Floresta
 761 Ombrófila Mista Em Campo Belo Do Sul , Sc. Cerne 10, 196–212.

762 Freitas, L., Sazima, M., 2006. Pollination Biology in a Tropical High-Altitude Grassland in Brazil:
 763 Interactions at the Community Level. *Ann. Missouri Bot. Gard.* 93, 465–516.

764 Garcia, M.J., De Oliveira, P.E., de Siqueira, E., Fernandes, R.S., 2004. A Holocene vegetational and
 765 climatic record from the Atlantic rainforest belt of coastal State of São Paulo, SE Brazil. *Rev.*
 766 *Palaeobot. Palynol.* 131, 181–199. <https://doi.org/10.1016/j.revpalbo.2004.03.007>

767 Gessert, S., Iriarte, J., Ríos, R.C., Behling, H., 2011. Late Holocene vegetation and environmental
 768 dynamics of the Araucaria forest region in Misiones Province, NE Argentina. *Rev. Palaeobot.*
 769 *Palynol.* 166, 29–37. <https://doi.org/10.1016/j.revpalbo.2011.04.006>

770 Gonçalves, E.T., Souza, A.F., 2014. Floristic variation in ecotonal areas: Patterns, determinants and
 771 biogeographic origins of subtropical forests in South America. *Austral Ecol.* 39, 122–134.
 772 <https://doi.org/10.1111/aec.12051>

773 Gosling, W.D., Mayle, F.E., Tate, N.J., Killeen, T.J., 2009. Differentiation between Neotropical
 774 rainforest, dry forest, and savannah ecosystems by their modern pollen spectra and
 775 implications for the fossil pollen record. *Rev. Palaeobot. Palynol.* 153, 70–85.
 776 <https://doi.org/10.1016/j.revpalbo.2008.06.007>

777 Gu, F., Zonneveld, K.A.F., Chiessi, C.M., Arz, H.W., Pätzold, J., Behling, H., 2017. Long-term
 778 vegetation, climate and ocean dynamics inferred from a 73,500 years old marine sediment
 779 core (GeoB2107-3) off southern Brazil. *Quat. Sci. Rev.* 172, 55–71.
 780 <https://doi.org/10.1016/j.quascirev.2017.06.028>

781 Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological Statistics Software Package for
 782 Education and Data Analysis. *Palaeontol. Electron.* 4, 1–9.
 783 <https://doi.org/10.1016/j.bcp.2008.05.025>

784 Haselhorst, D.S., Moreno, J.E., Punyasena, S.W., 2013. Variability within the 10-Year Pollen Rain of a
785 Seasonal Neotropical Forest and Its Implications for Paleoenvironmental and Phenological
786 Research. PLoS One 8. <https://doi.org/10.1371/journal.pone.0053485>

787 Haverroth, M., 1997. Kaingang, um estudo etnobotânico: o uso e a classificação das plantas na Área
788 Indígena Xapecó (oeste de SC). Universidade Federal de Santa Catarina.

789 Heineberg, M.R., 2014. Conhecimento e Uso das Plantas pelos Xokleng na TI Ibirama-Laklãnõ, Santa
790 Catarina. Universidade Federal de Santa Catarina.

791 Henry, J., 1964. Jungle people: A Kaingáng tribe of the highlands of Brazil, 2nd ed. Random House,
792 New York, NY.

793 Hermes, M.G., Köhler, A., 2006. The flower-visiting social wasps (Hymenoptera, Vespidae,
794 Polistinae) in two areas of Rio Grande do Sul State, southern Brazil. Rev. Bras. Entomol. 50,
795 268–274. <https://doi.org/10.1590/S0085-56262006000200008>

796 Hicks, S., 1985. Modern pollen deposition records from Kuusamo, Finland. Grana 24, 167–184.
797 <https://doi.org/10.1080/00173138509431005>

798 Hicks, S., 1977. Modern pollen rain in Finnish Lapland investigated by analysis of surface moss
799 samples. New Phytol. 78, 715–734. <https://doi.org/10.1111/j.1469-8137.1977.tb02176.x>

800 Higuchi, P., Silva, A.C. da, Sousa-Ferreira, T., Souza, S.T. de, Pereira-Gomes, J., Silva, K.M. da, Santos,
801 K.F. do, 2012. Floristic composition and phytogeography of the tree component of Araucaria
802 Forest fragments in southern Brazil. Brazilian J. Bot. 35, 145–157.
803 <https://doi.org/10.1590/S1806-99592012000200004>

804 Hill, M.O., Gauch, H.G., 1980. Detrended correspondence analysis: An improved ordination
805 technique. Vegetatio 42, 47–58. <https://doi.org/10.1007/BF00048870>

806 Hoffmann, K.D., 2011. Mito, história e alteridade entre os Xokleng. *Tellus* 11, 53–69.

807 Hooghiemstra, H., 1984. Vegetational and climatic history of the high plain of Bogota, Colombia: a
808 continuous record of the last 3.5 million years, *Dissertationes Botanicae*. J. Cramer.

809 Hueck, K., 1966. Die Wälder Südamerikas: Ökologie, Zusammensetzung und wirtschaftliche
810 Bedeutung. G. Fischer, Stuttgart.

811 Hueck, K., 1953. Distribuição e habitat natural do Pinheiro do Paraná (*Araucaria angustifolia*). *Bol.*
812 *da Fac. Filos. Ciências e Let. Univ. São Paulo*. 10, 5–24. [https://doi.org/10.11606/issn.2318-](https://doi.org/10.11606/issn.2318-5988.v10i1p5-24)
813 [5988.v10i1p5-24](https://doi.org/10.11606/issn.2318-5988.v10i1p5-24)

814 Imperatriz-Fonseca, V.L., Kleinert-Giovannini, A., Ramalho, M., 1989. Pollen Harvest by Eusocial
815 Bees in a Non-Natural Community in Brazil. *Source J. Trop. Ecol.* 5, 239–242.

816 Iriarte, J., Behling, H., 2007. The expansion of *Araucaria* forest in the southern Brazilian highlands
817 during the last 4000 years and its implications for the development of the Taquara/Itararé
818 Tradition. *Environ. Archaeol.* 12, 115–127. <https://doi.org/10.1179/174963107x226390>

819 Iriarte, J., DeBlasis, P., De Souza, J.G., Corteletti, R., 2016. Emergent Complexity, Changing
820 Landscapes, and Spheres of Interaction in Southeastern South America During the Middle and
821 Late Holocene. *J. Archaeol. Res.* 1–63. <https://doi.org/10.1007/s10814-016-9100-0>

822 Iriarte, J., Moehlecke Copé, S., Fradley, M., Lockhart, J.J., Gillam, J.C., 2013. Sacred landscapes of the
823 southern Brazilian highlands: Understanding southern proto-Je mound and enclosure
824 complexes. *J. Anthropol. Archaeol.* 32, 74–96. <https://doi.org/10.1016/j.jaa.2012.10.003>

825 Jeske-Pieruschka, V., Behling, H., 2012. Palaeoenvironmental history of the Sao Francisco de Paula
826 region in southern Brazil during the late Quaternary inferred from the Rincao das Cabritas
827 core. *The Holocene* 22, 1251–1262. <https://doi.org/10.1177/0959683611414930>

828 Jeske-Pieruschka, V., Fidelis, A., Bergamin, R.S., Vélez, E., Behling, H., 2010. Araucaria forest
 829 dynamics in relation to fire frequency in southern Brazil based on fossil and modern pollen
 830 data. *Rev. Palaeobot. Palynol.* 160, 53–65. <https://doi.org/10.1016/j.revpalbo.2010.01.005>

831 Jeske-Pieruschka, V., Pillar, V.D., de Oliveira, M.A.T., Behling, H., 2013. New insights into vegetation,
 832 climate and fire history of southern Brazil revealed by a 40,000 year environmental record
 833 from the State Park Serra do Tabuleiro. *Veg. Hist. Archaeobot.* 22, 299–314.
 834 <https://doi.org/10.1007/s00334-012-0382-y>

835 Juggins, S., 2016. C2 Software for Ecological and Palaeoecological Data Analysis and Visualisation.

836 Kent, M., Coker, P., 1992. Vegetation description and analysis : a practical approach. CRC Press.

837 Klein, R.M., 1978. Mapa fitogeográfico do Estado de Santa Catarina, Flora Ilustrada Catarinense.
 838 Herbário “Barbosa Rodrigues,” Itajai.

839 Klein, R.M., 1975. Southern Brazilian phytogeographic features and the probable influence of upper
 840 Quaternary climatic changes in the floristic distribution. *Bol. Parana. Geociências* 33, 67–88.

841 Lacerda, A.E.B., 2016. Conservation strategies for Araucaria Forests in Southern Brazil: assessing
 842 current and alternative approaches. *Biotropica* 48, 537–544.
 843 <https://doi.org/10.1111/btp.12317>

844 Lauterjung, M.B., Bernardi, A.P., Montagna, T., Candido-Ribeiro, R., da Costa, N.C.F., Mantovani, A.,
 845 dos Reis, M.S., 2018. Phylogeography of Brazilian pine (*Araucaria angustifolia*): integrative
 846 evidence for pre-Columbian anthropogenic dispersal. *Tree Genet. Genomes* 14, 36.
 847 <https://doi.org/10.1007/s11295-018-1250-4>

848 Ledru, M.-P., Mourguiart, P., Riccomini, C., 2009. Related changes in biodiversity, insolation and
 849 climate in the Atlantic rainforest since the last interglacial. *Palaeogeogr. Palaeoclimatol.*

850 Palaeoecol. 271, 140–152. <https://doi.org/10.1016/j.palaeo.2008.10.008>

851 Legendre, P., Legendre, L., Legendre, L., Legendre, P., 2012. Numerical ecology. Elsevier.

852 Leite, P.F., Klein, R.M., 1990. Vegetação, in: Mesquita, O.V. (Ed.), Geografia Do Brasil — Região Sul
853 (Vol. 2). Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro, pp. 113–150.

854 Leonhardt, A., Lorscheitter, M.L., 2010. The last 25,000 years in the Eastern Plateau of Southern
855 Brazil according to Alpes de Sao Francisco record. J. South Am. Earth Sci. 29, 454–463.
856 <https://doi.org/10.1016/j.jsames.2009.09.003>

857 Levis, C., Costa, F.R.C., Bongers, F., Peña-Claros, M., Clement, C.R., Junqueira, A.B., Neves, E.G.,
858 Tamanaha, E.K., Figueiredo, F.O.G., Salomão, R.P., Castilho, C. V, Magnusson, W.E., Phillips,
859 O.L., Guevara, J.E., Sabatier, D., Molino, J.-F., López, D.C., Mendoza, A.M., Pitman, N.C.A.,
860 Duque, A., Vargas, P.N., Zartman, C.E., Vasquez, R., Andrade, A., Camargo, J.L., Feldpausch,
861 T.R., Laurance, S.G.W., Laurance, W.F., Killeen, T.J., Nascimento, H.E.M., Montero, J.C.,
862 Mostacedo, B., Amaral, I.L., Guimarães Vieira, I.C., Brienens, R., Castellanos, H., Terborgh, J.,
863 Carim, M. de J.V., Guimarães, J.R. da S., Coelho, L.D.S., Matos, F.D.D.A., Wittmann, F.,
864 Mogollón, H.F., Damasco, G., Dávila, N., García-Villacorta, R., Coronado, E.N.H., Emilio, T.,
865 Filho, D. de A.L., Schietti, J., Souza, P., Targhetta, N., Comiskey, J.A., Marimon, B.S., Marimon,
866 B.-H., Neill, D., Alonso, A., Arroyo, L., Carvalho, F.A., de Souza, F.C., Dallmeier, F., Pansonato,
867 M.P., Duivenvoorden, J.F., Fine, P.V.A., Stevenson, P.R., Araujo-Murakami, A., Aymard C., G.A.,
868 Baraloto, C., do Amaral, D.D., Engel, J., Henkel, T.W., Maas, P., Petronelli, P., Revilla, J.D.C.,
869 Stropp, J., Daly, D., Gribel, R., Paredes, M.R., Silveira, M., Thomas-Caesar, R., Baker, T.R., da
870 Silva, N.F., Ferreira, L. V., Peres, C.A., Silman, M.R., Cerón, C., Valverde, F.C., Di Fiore, A.,
871 Jimenez, E.M., Mora, M.C.P., Toledo, M., Barbosa, E.M., Bonates, L.C. de M., Arboleda, N.C.,
872 Farias, E. de S., Fuentes, A., Guillaumet, J.-L., Jørgensen, P.M., Malhi, Y., de Andrade Miranda,

873 I.P., Phillips, J.F., Prieto, A., Rudas, A., Ruschel, A.R., Silva, N., von Hildebrand, P., Vos, V.A.,
 874 Zent, E.L., Zent, S., Cintra, B.B.L., Nascimento, M.T., Oliveira, A.A., Ramirez-Angulo, H., Ramos,
 875 J.F., Rivas, G., Schöngart, J., Sierra, R., Tirado, M., van der Heijden, G., Torre, E. V., Wang, O.,
 876 Young, K.R., Baider, C., Cano, A., Farfan-Rios, W., Ferreira, C., Hoffman, B., Mendoza, C.,
 877 Mesones, I., Torres-Lezama, A., Medina, M.N.U., van Andel, T.R., Villarroel, D., Zagt, R.,
 878 Alexiades, M.N., Balslev, H., Garcia-Cabrera, K., Gonzales, T., Hernandez, L., Huamantupa-
 879 Chuquimaco, I., Manzatto, A.G., Milliken, W., Cuenca, W.P., Pansini, S., Pauletto, D., Arevalo,
 880 F.R., Reis, N.F.C., Sampaio, A.F., Giraldo, L.E.U., Sandoval, E.H.V., Gamarra, L.V., Vela, C.I.A., ter
 881 Steege, H., 2017. Persistent effects of pre-Columbian plant domestication on Amazonian forest
 882 composition. *Science* (80-.). 355, 925–931. <https://doi.org/10.1126/science.aal0157>
 883 Levis, C., Flores, B.M., Moreira, P.A., Luize, B.G., Alves, R.P., Franco-Moraes, J., Lins, J., Konings, E.,
 884 Peña-Claros, M., Bongers, F., Costa, F.R.C., Clement, C.R., 2018. How People Domesticated
 885 Amazonian Forests. *Front. Ecol. Evol.* 5, 1–21. <https://doi.org/10.3389/fevo.2017.00171>
 886 López-Sáez, J.A., Alba-Sánchez, F., López-Merino, L., Pérez-Díaz, S., 2010. Modern pollen analysis: a
 887 reliable tool for discriminating *Quercus rotundifolia* communities in Central Spain.
 888 *Phytocoenologia* 40, 57–72. <https://doi.org/10.1127/0340-269X/2010/0040-0430>
 889 Loponte, D., Carbonera, M., Corriale, M.J., Acosta, A., 2016. Horticulturists and oxygen ecozones in
 890 the tropical and subtropical forests of Southeast South America. *Environ. Archaeol.* 22, 1–21.
 891 <https://doi.org/10.1080/14614103.2016.1211382>
 892 Maçaneiro, J.P. DE, Gasper, A.L. DE, Galvão, F., Schorn, L.A., 2018. Dispersion and aggregation
 893 patterns of tree species in Araucaria Forest, Southern Brazil. *An. Acad. Bras. Cienc.* 90, 2397–
 894 2408. <https://doi.org/10.1590/0001-3765201820170150>

895 McMichael, C.N.H., Matthews-Bird, F., Farfan-Rios, W., Feeley, K.J., 2017. Ancient human
 896 disturbances may be skewing our understanding of Amazonian forests. *Proc. Natl. Acad. Sci.*
 897 114, 522–527. <https://doi.org/10.1073/pnas.1614577114>

898 Mello, A.J.M., Peroni, N., 2015. Cultural landscapes of the Araucaria Forests in the northern plateau
 899 of Santa Catarina, Brazil. *J. Ethnobiol. Ethnomed.* 11, 51. [https://doi.org/10.1186/s13002-015-](https://doi.org/10.1186/s13002-015-0039-x)
 900 0039-x

901 Métraux, A., 1946. The Caingang, in: Steward, J.H. (Ed.), *Handbook of South American Indians. The*
 902 *Marginal Tribes*. Smithsonian Institution, Washington (D.C.), pp. 445–475.

903 Meyer, L., Sevegnani, L., de Gasper, A.L., Schorn, L.A., Vibrans, A.C., Lingner, D.V., Sobral, M.G.,
 904 Klemz, G., Schmitt, R., Anastacio Jr., C., Brogni, E., 2013. Fitossociologia do componente
 905 arbóreo/arbustivo da Floresta Ombrófila Mista no Estado de Santa Catarina, in: Vibrans, A.C.,
 906 Sevegnani, L., Gasper, A.L. De, Lingner, D.V. (Eds.), *Inventário Florístico Florestal de Santa*
 907 *Catarina Volume III - Floresta Ombrófila Mista*. Edifurb, Blumenau, pp. 157–189.

908 Midgley, J., 1989. Pollen dispersal distances for a conifer canopy species in the Knysna Forest. *South*
 909 *African J. Bot.* 55, 662–663.

910 Montade, V., Diogo, I.J.S., Bremond, L., Favier, C., Costa, I.R. da, Ledru, M.P., Paradis, L., Martins,
 911 E.S.P.R., Burte, J., Silva, F.H.M., Verola, C.F., 2016. Pollen-based characterization of montane
 912 forest types in north-eastern Brazil. *Rev. Palaeobot. Palynol.* 234, 147–158.
 913 <https://doi.org/10.1016/j.revpalbo.2016.07.003>

914 Montalva, J., Dudley, L., Arroyo, M.K., Retamales, H., Abrahamovich, A.H., 2011. Geographic
 915 distribution and associated flora of native and introduced bumble bees (*Bombus* spp.) in Chile.
 916 *J. Apic. Res.* 50, 11–21.

917 Mrotzek, A., Couwenberg, J., Theuerkauf, M., Joosten, H., 2017. MARCO POLO - A new and simple
 918 tool for pollen-based stand-scale vegetation reconstruction. *The Holocene* 27, 321–330.
 919 <https://doi.org/10.1177/0959683616660171>

920 Müller, S.C., Overbeck, G.E., Blanco, C.C., de Oliveira, J.M., Pillar, V.D., 2012. South Brazilian Forest-
 921 Grassland Ecotones: Dynamics Affected by Climate, Disturbance, and Woody Species Traits, in:
 922 Myser, R.W. (Ed.), *Ecotones Between Forest and Grassland*. Springer New York, New York, NY,
 923 pp. 167–187. https://doi.org/10.1007/978-1-4614-3797-0_7

924 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J., 2000. Biodiversity
 925 hotspots for conservation priorities. *Nature* 403, 853–858. <https://doi.org/10.1038/35002501>

926 Negash, L., 2003. In situ fertility decline and provenance differences in the East African Yellow
 927 Wood (*Podocarpus falcatus*) measured through in vitro seed germination. *For. Ecol. Manage.*
 928 174, 127–138. [https://doi.org/10.1016/S0378-1127\(02\)00034-8](https://doi.org/10.1016/S0378-1127(02)00034-8)

929 Neves, D.M., Dexter, K.G., Pennington, R.T., Valente, A.S.M., Bueno, M.L., Eisenlohr, P. V., Fontes,
 930 M.A.L., Miranda, P.L.S., Moreira, S.N., Rezende, V.L., Saiter, F.Z., Oliveira-Filho, A.T., 2017.
 931 Dissecting a biodiversity hotspot: The importance of environmentally marginal habitats in the
 932 Atlantic Forest Domain of South America. *Divers. Distrib.* 23, 898–909.
 933 <https://doi.org/10.1111/ddi.12581>

934 Noelli, F.S., 2000. A ocupação humana na Região Sul do Brasil: Arqueologia, debates e perspectivas
 935 1872-2000. *Rev. da Univ. da São Paulo* 44, 218–269.

936 Noelli, F.S., De Souza, J.G., 2017. Novas perspectivas para a cartografia arqueológica Jê no Brasil
 937 meridional. *Bol. do Mus. Para. Emilio Goeldi Ciencias Humanas*.
 938 <https://doi.org/10.1590/1981.81222017000100004>

939 Oksanen, J., Minchin, P.R., 1997. Instability of ordination results under changes in input data order:
 940 explanations and remedies. *J. Veg. Sci.* 8, 447–454. <https://doi.org/10.2307/3237336>

941 Oliveira-Filho, A.T., Budke, J.C., Jarenkow, J.A., Eisenlohr, P. V, Neves, D.R.M., 2014. Delving into the
 942 variations in tree species composition and richness across South American subtropical Atlantic
 943 and Pampean forests. *J. Plant Ecol.* 8, 242–260. <https://doi.org/10.1093/jpe/rtt058>

944 Otegui, M., Cocucci, A., 1999. Flower morphology and biology of *Myrsine laetevirens*, structural and
 945 evolutionary implications of anemophily in Myrsinaceae. *Nord. J. Bot.* 19, 71–85.
 946 <https://doi.org/10.1111/j.1756-1051.1999.tb01904.x>

947 Pardoe, H.S., 2006. Surface pollen deposition on glacier forelands in southern Norway I: local
 948 patterns of representation and source area at Storbreen, Jotunheimen. *The Holocene* 16,
 949 1149–1161. <https://doi.org/10.1177/0959683606069422>

950 Pardoe, H.S., Giesecke, T., van der Knaap, W.O., Svitavská-Svobodová, H., Kvavadze, E. V.,
 951 Panajiotidis, S., Gerasimidis, A., Pidek, I.A., Zimny, M., Święta-Musznicka, J., Latałowa, M.,
 952 Noryśkiewicz, A.M., Bozilova, E., Tonkov, S., Filipova-Marinova, M. V., van Leeuwen, J.F.N.,
 953 Kalniņa, L., 2010. Comparing pollen spectra from modified Tauber traps and moss samples:
 954 examples from a selection of woodlands across Europe. *Veg. Hist. Archaeobot.* 19, 271–283.
 955 <https://doi.org/10.1007/s00334-010-0258-y>

956 Parsons, R.W., Prentice, I.C., 1981. Statistical approaches to R-values and the pollen—vegetation
 957 relationship. *Rev. Palaeobot. Palynol.* 32, 127–152. [https://doi.org/10.1016/0034-](https://doi.org/10.1016/0034-6667(81)90001-4)
 958 [6667\(81\)90001-4](https://doi.org/10.1016/0034-6667(81)90001-4)

959 Peterson, R.H., Gale, D., 1991. Fish species associations in riffle habitat of streams of varying size
 960 and acidity in New Brunswick and Nova Scotia. *J. Fish Biol.* 38, 859–871.

961 <https://doi.org/10.1111/j.1095-8649.1991.tb03626.x>

962 R Core Team, 2017. R: a Language and Environment for Statistical Computing.

963 Reis, M.S., Montagna, T., Mattos, A.G., Filippon, S., Ladio, A.H., Marques, A. da C., Zechini, A.A.,
 964 Peroni, N., Mantovani, A., 2018. Domesticated Landscapes in Araucaria Forests, Southern
 965 Brazil: A Multispecies Local Conservation-by-Use System. *Front. Ecol. Evol.* 6, 11.
 966 <https://doi.org/10.3389/fevo.2018.00011>

967 Rezende, V.L., Eisenlohr, P. V., Vibrans, A.C., de Oliveira-Filho, A.T., 2015. Humidity, low
 968 temperature extremes, and space influence floristic variation across an insightful gradient in
 969 the Subtropical Atlantic Forest. *Plant Ecol.* 216, 759–774. [https://doi.org/10.1007/s11258-015-](https://doi.org/10.1007/s11258-015-0465-9)
 970 [0465-9](https://doi.org/10.1007/s11258-015-0465-9)

971 Ribeiro, M.C., Metzger, J.P., Martensen, A.C., Ponzoni, F.J., Hirota, M.M., 2009. The Brazilian
 972 Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for
 973 conservation. *Biol. Conserv.* 142, 1141–1153. <https://doi.org/10.1016/j.biocon.2009.02.021>

974 Robinson, M., De Souza, J.G., Maezumi, S.Y., Cárdenas, M., Pessenda, L., Prufer, K., Corteletti, R.,
 975 Scunderlick, D., Mayle, F.E., De Blasis, P., Iriarte, J., 2018. Uncoupling human and climate
 976 drivers of late Holocene vegetation change in southern Brazil. *Sci. Rep.* 8, 7800.
 977 <https://doi.org/10.1038/s41598-018-24429-5>

978 Robinson, M., Iriarte, J., De Souza, J.G., Marozzi, O., Scheel-Ybert, R., 2017. Moiety specific wood
 979 selection in funerary ritual for the southern proto-Jê. *J. Archaeol. Sci. Reports* 11, 237–244.
 980 <https://doi.org/10.1016/j.jasrep.2016.11.047>

981 Rodrigues, J.M., Behling, H., Giesecke, T., 2016a. Holocene dynamics of vegetation change in
 982 southern and southeastern Brazil is consistent with climate forcing. *Quat. Sci. Rev.* 146, 54–65.

<https://doi.org/10.1016/j.quascirev.2016.06.011>

Rodrigues, J.M., Behling, H., Giesecke, T., 2016b. Differentiating vegetation types from eastern South American ecosystems based on modern and subfossil pollen samples: evaluating modern analogues. *Veg. Hist. Archaeobot.* 25, 387–403. <https://doi.org/10.1007/s00334-016-0558-y>

Roubik, D.W., Moreno, J.E., 1991. Pollen and spores of Barro Colorado Island (Panama), Monographs in Systematic Botany. Missouri Botanical Garden, St. Louis, MO, USA.

Scherer, C., Lorscheitter, M.L., 2014. Vegetation dynamics in the southern Brazilian highlands during the last millennia and the role of bogs in Araucaria forest formation. *Quat. Int.* 325, 3–12. <https://doi.org/10.1016/j.quaint.2014.01.010>

Schorn, L.A., de Gasper, A.L., Meyer, L., Vibrans, A.C., 2012. Síntese da estrutura dos remanescentes florestais em Santa Catarina, in: Vibrans, A.C., Sevegnani, L., de Gasper, A.L., Lingner, D.V. (Eds.), *Inventário Florístico Florestal de Santa Catarina Volume I - Diversidade e Conservação Dos Remanescentes Florestais*. Edifurb, Blumenau, pp. 125–140.

Sevegnani, L., Uhlmann, A., de Gasper, A.L., Vibrans, A.C., dos Santos, A.S., Verdi, M., drev, 2013. Estádios sucessionais da Floresta Ombrófila Mista em Santa Catarina, in: Vibrans, A.C., Sevegnani, L., de Gaspar, A.L., Lingner, D.V. (Eds.), *Inventário Florístico Florestal de Santa Catarina Volume III - Floresta Ombrófila Mista*. Edifurb, Blumenau, p. 440.

Sevegnani, L., Uhlmann, A., Gasper, A.L. de, Meyer, L., Vibrans, A.C., 2016. Climate affects the structure of mixed rain forest in southern sector of Atlantic domain in Brazil. *Acta Oecologica* 77, 109–117. <https://doi.org/10.1016/j.actao.2016.10.002>

Shepard, G., Ramirez, H., 2011. “Made in Brazil”: Human Dispersal of the Brazil Nut (*Bertholletia*

1005 excelsa, Lecythidaceae) in Ancient Amazonia. Econ. Bot. 65, 44–65.

1006 <https://doi.org/10.1007/s12231-011-9151-6>

1007 Silva, S.B. da, 2002. Dualismo e cosmologia Kaingang: o xamã e o domínio da floresta. Horizontes

1008 Antropológicos 8, 189–209. <https://doi.org/10.1590/S0104-71832002000200009>

1009 Silva, L.C.R., Anand, M., 2011. Mechanisms of Araucaria (Atlantic) Forest Expansion into Southern

1010 Brazilian Grasslands. Ecosystems 14, 1354–1371. <https://doi.org/10.1007/s10021-011-9486-y>

1011 Souza, L.A. de, Moscheta, I.S., 2000. Morfo-anatomia e aspectos da biologia floral de Nectandra

1012 megapotamica (Spreng.) Mez (Lauraceae). Acta Bot. Brasilica 14, 15–25.

1013 <https://doi.org/10.1590/S0102-33062000000100003>

1014 Sugita, S., 2007a. Theory of quantitative reconstruction of vegetation I: pollen from large sites

1015 REVEALS regional vegetation composition. The Holocene 17, 229–241.

1016 <https://doi.org/10.1177/0959683607075837>

1017 Sugita, S., 2007b. Theory of quantitative reconstruction of vegetation II: all you need is LOVE. The

1018 Holocene 17, 243–257. <https://doi.org/10.1177/0959683607075838>

1019 Thomas, P., 2013. Araucaria angustifolia [WWW Document]. IUCN Red List Threat. Species.

1020 <https://doi.org/http://dx.doi.org/10.2305/IUCN.UK.2013-1.RLTS.T32975A2829141.en>

1021 Tomlinson, P.B., 1974. Breeding Mechanisms in Trees Native to Tropical Florida - a Morphological

1022 Assessment. J. Arnold Arbor. 55, 269–290. <https://doi.org/10.2307/43781937>

1023 Tonkov, S., Hicks, S., Bozilova, E., Atanassova, J., 2001. Pollen monitoring in the central Rila

1024 Mountains, Southwestern Bulgaria: comparisons between pollen traps and surface samples for

1025 the period 1993–1999. Rev. Palaeobot. Palynol. 117, 167–182. [https://doi.org/10.1016/S0034-](https://doi.org/10.1016/S0034-6667(01)00085-9)

1026 6667(01)00085-9

- 1027 Urban, G., 1985. Interpretations of Inter-Cultural Contact: The Shokleng and Brazilian National
1028 Society 1914-1915. *Ethnohistory* 32, 224–244. <https://doi.org/10.2307/481922>
- 1029 Vibrans, A.C., Sevegnani, L., Lingner, D.V., de Gasper, A.L., Sabbagh, S., 2010. Inventário florístico
1030 florestal de Santa Catarina (IFFSC): aspectos metodológicos e operacionais. *Pesqui. Florest.*
1031 *Bras.* 30, 291–302. <https://doi.org/10.4336/2010.pfb.64.291>
- 1032 Viera, F. V., Fonseca, R.S., Araújo, L.M. de, 2012. Floração, Polinização e Sistemas Reprodutivos em
1033 Florestas Tropicais, in: Martins, S.V. (Ed.), *Ecologia de Florestas Tropicais Do Brasil*.
1034 Universidade Federal de Viçosa, Viçosa, pp. 53–84.
- 1035 Whitehead, D.R., 1983. Wind pollination: some ecological and evolutionary perspectives, in: Real, L.
1036 (Ed.), *Pollination Biology*. Academic Press, Inc., Orlando, pp. 97–108.
- 1037 Whitney, B.S., Smallman, T.L., Mitchard, E.T., Carson, J.F., Mayle, F.E., Bunting, M.J., 2019.
1038 Constraining pollen-based estimates of forest cover in the Amazon: A simulation approach.
1039 *The Holocene* 29, 262–270. <https://doi.org/10.1177/0959683618810394>
- 1040 Wilms, W., Wendel, L., Zillikens, A., Blochtein, B., Engels, W., 1997. Bees and Other Insects
1041 Recorded on Flowering Trees in a Subtropical Araucaria Forest in Southern Brazil. *Stud.*
1042 *Neotrop. Fauna Environ.* 32, 220–226.
1043 <https://doi.org/10.1080/01650521.1997.11432425>
- 1044 Wilson, V.R., Owens, J.N., 1999. The Reproductive Biology of Totara (*Podocarpus totara*)
1045 (*Podocarpaceae*). *Ann. Bot.* 83, 401–411.
- 1046 Wrege, M.S., Higa, R.C. V., Miranda Britez, R., Cordeiro Garrastazu, M., de Sousa, V.A., Caramori,
1047 P.H., Radin, B., Braga, H.J., 2009. Climate Change and Conservation of *Araucaria angustifolia* in
1048 Brazil. *Unasylva* 60, 30–33.

1049 Zeller, R.H., 2010. PLANO DE MANEJO: RESERVA PARTICULAR DO PATRIMÔNIO NATURAL EMILIO
1050 EINSFELD FILHO, SANTA CATARINA. Campo Belo do Sul.

1051

1052

1053

1054 **Figures**

1055 Figure 1: Map of forest plot locations; numbers refer to plot codes (see Table 1 below). Plot colours:

1056 Cyan: riverine; Olive: slope; Black: rock outcrop; Purple: open understorey; Orange: closed

1057 understorey; Red: disturbed.




Figure 2: Relative abundance (%) of selected tree species in the vegetation survey. Species included in the cluster analysis and DCA are indicated with an asterisk.






Figure 3: Relative Coverage Values (RCVs, %) of selected tree species in the vegetation survey. 'Ave. (all)' = sum of RCVs/total number of plots; 'Ave. (present)' = sum of RCVs/number of plots in which taxon was present.



1069 Figure 4: Relative abundance (%) of selected pollen types. Taxa included in the cluster analysis and
1070 DCA are indicated with an asterisk



1078 Figure 6: p/v values of key genera of *Araucaria* forest in this study. *Clethra* has an infinitely high p/v
1079 value; faint sections in the bars of *Lamanonia* and *Lithraea* reflect alternate calculations of their p/v
1080 values (see section 3.4).

